Technical Support Document: -Technical Update of the Social Cost of Carbon for Regulatory Impact Analysis -Under Executive Order 12866 -

Interagency Working Group on Social Cost of Carbon, United States Government

With participation by

Council of Economic Advisers Council on Environmental Quality Department of Agriculture Department of Commerce Department of Energy Department of Transportation Domestic Policy Council Environmental Protection Agency National Economic Council Office of Management and Budget Office of Science and Technology Policy Department of the Treasury

May 2013

Revised November 2013 See Appendix B for Details on Revision

Executive Summary

Under Executive Order 12866, agencies are required, to the extent permitted by law, "to assess both the costs and the benefits of the intended regulation and, recognizing that some costs and benefits are difficult to quantify, propose or adopt a regulation only upon a reasoned determination that the benefits of the intended regulation justify its costs." The purpose of the "social cost of carbon" (SCC) estimates presented here is to allow agencies to incorporate the social benefits of reducing carbon dioxide (CO₂) emissions into cost-benefit analyses of regulatory actions that impact cumulative global emissions. The SCC is an estimate of the monetized damages associated with an incremental increase in carbon emissions in a given year. It is intended to include (but is not limited to) changes in net agricultural productivity, human health, property damages from increased flood risk, and the value of ecosystem services due to climate change.

The interagency process that developed the original U.S. government's SCC estimates is described in the 2010 interagency technical support document (TSD) (Interagency Working Group on Social Cost of Carbon 2010). Through that process the interagency group selected four SCC values for use in regulatory analyses. Three values are based on the average SCC from three integrated assessment models (IAMs), at discount rates of 2.5, 3, and 5 percent. The fourth value, which represents the 95th percentile SCC estimate across all three models at a 3 percent discount rate, is included to represent higher-than-expected impacts from temperature change further out in the tails of the SCC distribution.

While acknowledging the continued limitations of the approach taken by the interagency group in 2010, this document provides an update of the SCC estimates based on new versions of each IAM (DICE, PAGE, and FUND). It does not revisit other interagency modeling decisions (e.g., with regard to the discount rate, reference case socioeconomic and emission scenarios, or equilibrium climate sensitivity). Improvements in the way damages are modeled are confined to those that have been incorporated into the latest versions of the models by the developers themselves in the peer-reviewed literature.

The SCC estimates using the updated versions of the models are higher than those reported in the 2010 TSD. By way of comparison, the four 2020 SCC estimates reported in the 2010 TSD were \$7, \$26, \$42 and \$81 (2007\$). The corresponding four updated SCC estimates for 2020 are \$12, \$43, \$64, and \$128 (2007\$). The model updates that are relevant to the SCC estimates include: an explicit representation of sea level rise damages in the DICE and PAGE models; updated adaptation assumptions, revisions to ensure damages are constrained by GDP, updated regional scaling of damages, and a revised treatment of potentially abrupt shifts in climate damages in the PAGE model; an updated carbon cycle in the DICE model; and updated damage functions for sea level rise impacts, the agricultural sector, and reduced space heating requirements, as well as changes to the transient response of temperature to the buildup of GHG concentrations and the inclusion of indirect effects of methane emissions in the FUND model. The SCC estimates vary by year, and the following table summarizes the revised SCC estimates from 2010 through 2050.

Discount Rate Year	5.0% Avg	3.0% Avg	2.5% Avg	3.0% 95th
2010	11	32	51	89
2015	11	37	57	109
2020	12	43	64	128
2025	14	47	69	143
2030	16	52	75	159
2035	19	56	80	175
2040	21	61	86	191
2045	24	66	92	206
2050	26	71	97	220

Revised Social Cost of CO₂, 2010 – 2050 (in 2007 dollars per metric ton of CO₂)

I. Purpose

The purpose of this document is to update the schedule of social cost of carbon (SCC) estimates from the 2010 interagency technical support document (TSD) (Interagency Working Group on Social Cost of Carbon 2010).¹ E.O. 13563 commits the Administration to regulatory decision making "based on the best available science."² Additionally, the interagency group recommended in 2010 that the SCC estimates be revisited on a regular basis or as model updates that reflect the growing body of scientific and economic knowledge become available.³ New versions of the three integrated assessment models used by the U.S. government to estimate the SCC (DICE, FUND, and PAGE), are now available and have been published in the peer reviewed literature. While acknowledging the continued limitations of the approach taken by the interagency group in 2010 (documented in the original 2010 TSD), this document provides an update of the SCC estimates based on the latest peer-reviewed version of the models, replacing model versions that were developed up to ten years ago in a rapidly evolving field. It does not revisit other assumptions with regard to the discount rate, reference case socioeconomic and emission scenarios, or equilibrium climate sensitivity. Improvements in the way damages are modeled are confined to those that have been incorporated into the latest versions of the models by the developers themselves in the peer-reviewed literature. The agencies participating in the interagency working group continue to investigate potential improvements to the way in which economic damages associated with changes in CO₂ emissions are quantified.

Section II summarizes the major updates relevant to SCC estimation that are contained in the new versions of the integrated assessment models released since the 2010 interagency report. Section III presents the updated schedule of SCC estimates for 2010 – 2050 based on these versions of the models. Section IV provides a discussion of other model limitations and research gaps.

II. Summary of Model Updates

This section briefly summarizes changes to the most recent versions of the three integrated assessment models (IAMs) used by the interagency group in 2010. We focus on describing those model updates that are relevant to estimating the social cost of carbon, as summarized in Table 1. For example, both the DICE and PAGE models now include an explicit representation of sea level rise damages. Other revisions to PAGE include: updated adaptation assumptions, revisions to ensure damages are constrained by GDP, updated regional scaling of damages, and a revised treatment of potentially abrupt shifts in climate damages. The DICE model's simple carbon cycle has been updated to be more consistent with a more complex climate model. The FUND model includes updated damage functions for sea level rise impacts, the agricultural sector, and reduced space heating requirements, as well as changes to the transient response of temperature to the buildup of GHG concentrations and the inclusion of indirect effects of

¹ In this document, we present all values of the SCC as the cost per metric ton of CO₂ emissions. Alternatively, one could report the SCC as the cost per metric ton of carbon emissions. The multiplier for translating between mass of CO₂ and the mass of carbon is 3.67 (the molecular weight of CO₂ divided by the molecular weight of carbon = 44/12 = 3.67).

² http://www.whitehouse.gov/sites/default/files/omb/inforeg/eo12866/eo13563_01182011.pdf

³ See p. 1, 3, 4, 29, and 33 (Interagency Working Group on Social Cost of Carbon 2010).

methane emissions. Changes made to parts of the models that are superseded by the interagency working group's modeling assumptions – regarding equilibrium climate sensitivity, discounting, and socioeconomic variables – are not discussed here but can be found in the references provided in each section below.

IAM	Version used in 2010 Interagency Analysis	New Version	Key changes relevant to interagency SCC
DICE	2007	2010	Updated calibration of the carbon cycle model and explicit representation of sea level rise (SLR) and associated damages.
FUND	3.5 (2009)	3.8 (2012)	Updated damage functions for space heating, SLR, agricultural impacts, changes to transient response of temperature to buildup of GHG concentrations, and inclusion of indirect climate effects of methane.
PAGE	2002	2009	Explicit representation of SLR damages, revisions to damage function to ensure damages do not exceed 100% of GDP, change in regional scaling of damages, revised treatment of potential abrupt damages, and updated adaptation assumptions.

Table 1: Summary of Key Model Revisions Relevant to the Interagency SCC

A. DICE

DICE 2010 includes a number of changes over the previous 2007 version used in the 2010 interagency report. The model changes that are relevant for the SCC estimates developed by the interagency working group include: 1) updated parameter values for the carbon cycle model, 2) an explicit representation of sea level dynamics, and 3) a re-calibrated damage function that includes an explicit representation of economic damages from sea level rise. Changes were also made to other parts of the DICE model—including the equilibrium climate sensitivity parameter, the rate of change of total factor productivity, and the elasticity of the marginal utility of consumption—but these components of DICE are superseded by the interagency working group's assumptions and so will not be discussed here. More details on DICE2007 can be found in Nordhaus (2008) and on DICE2010 in Nordhaus (2010). The DICE2010 model and documentation is also available for download from the homepage of William Nordhaus.

Carbon Cycle Parameters

DICE uses a three-box model of carbon stocks and flows to represent the accumulation and transfer of carbon among the atmosphere, the shallow ocean and terrestrial biosphere, and the deep ocean. These parameters are "calibrated to match the carbon cycle in the Model for the Assessment of Greenhouse

Gas Induced Climate Change (MAGICC)" (Nordhaus 2008 p 44).⁴ Carbon cycle transfer coefficient values in DICE2010 are based on re-calibration of the model to match the newer 2009 version of MAGICC (Nordhaus 2010 p 2). For example, in DICE2010, in each decade, 12 percent of the carbon in the atmosphere is transferred to the shallow ocean, 4.7 percent of the carbon in the shallow ocean is transferred to the atmosphere, 94.8 percent remains in the shallow ocean, and 0.5 percent is transferred to the deep ocean. For comparison, in DICE 2007, 18.9 percent of the carbon in the atmosphere is transferred to the shallow ocean each decade, 9.7 percent of the carbon in the shallow ocean is transferred to the atmosphere, 85.3 percent remains in the shallow ocean, and 5 percent is transferred to the deep ocean.

The implication of these changes for DICE2010 is in general a weakening of the ocean as a carbon sink and therefore a higher concentration of carbon in the atmosphere than in DICE2007, for a given path of emissions. All else equal, these changes will generally increase the level of warming and therefore the SCC estimates in DICE2010 relative to those from DICE2007.

Sea Level Dynamics

A new feature of DICE2010 is an explicit representation of the dynamics of the global average sea level anomaly to be used in the updated damage function (discussed below). This section contains a brief description of the sea level rise (SLR) module; a more detailed description can be found on the model developer's website.⁵ The average global sea level anomaly is modeled as the sum of four terms that represent contributions from: 1) thermal expansion of the oceans, 2) melting of glaciers and small ice caps, 3) melting of the Greenland ice sheet, and 4) melting of the Antarctic ice sheet.

The parameters of the four components of the SLR module are calibrated to match consensus results from the IPCC's Fourth Assessment Report (AR4).⁶ The rise in sea level from thermal expansion in each time period (decade) is 2 percent of the difference between the sea level in the previous period and the long run equilibrium sea level, which is 0.5 meters per degree Celsius (°C) above the average global temperature in 1900. The rise in sea level from the melting of glaciers and small ice caps occurs at a rate of 0.008 meters per decade per °C above the average global temperature in 1900.

The contribution to sea level rise from melting of the Greenland ice sheet is more complex. The equilibrium contribution to SLR is 0 meters for temperature anomalies less than 1 °C and increases linearly from 0 meters to a maximum of 7.3 meters for temperature anomalies between 1 °C and 3.5 °C. The contribution to SLR in each period is proportional to the difference between the previous period's sea level anomaly and the equilibrium sea level anomaly, where the constant of proportionality increases with the temperature anomaly in the current period.

⁴ MAGICC is a simple climate model initially developed by the U.S. National Center for Atmospheric Research that has been used heavily by the Intergovernmental Panel on Climate Change (IPCC) to emulate projections from more sophisticated state of the art earth system simulation models (Randall et al. 2007).

⁵ Documentation on the new sea level rise module of DICE is available on William Nordhaus' website at: http://nordhaus.econ.yale.edu/documents/SLR_021910.pdf.

⁶ For a review of post-IPCC AR4 research on sea level rise, see Nicholls et al. (2011) and NAS (2011).

The contribution to SLR from the melting of the Antarctic ice sheet is -0.001 meters per decade when the temperature anomaly is below 3 °C and increases linearly between 3 °C and 6 °C to a maximum rate of 0.025 meters per decade at a temperature anomaly of 6 °C.

Re-calibrated Damage Function

Economic damages from climate change in the DICE model are represented by a fractional loss of gross economic output in each period. A portion of the remaining economic output in each period (net of climate change damages) is consumed and the remainder is invested in the physical capital stock to support future economic production, so each period's climate damages will reduce consumption in that period and in all future periods due to the lost investment. The fraction of output in each period that is lost due to climate change impacts is represented as one minus a fraction, which is one divided by a quadratic function of the temperature anomaly, producing a sigmoid ("S"-shaped) function.⁷ The loss function in DICE2010 has been expanded by adding a quadratic function of SLR to the quadratic function of temperature. In DICE2010 the temperature anomaly coefficients have been recalibrated to avoid double-counting damages from sea level rise that were implicitly included in these parameters in DICE2007.

The aggregate damages in DICE2010 are illustrated by Nordhaus (2010 p 3), who notes that "...damages in the uncontrolled (baseline) [i.e., reference] case ... in 2095 are \$12 trillion, or 2.8 percent of global output, for a global temperature increase of 3.4 °C above 1900 levels." This compares to a loss of 3.2 percent of global output at 3.4 °C in DICE2007. However, in DICE2010, annual damages are lower in most of the early periods of the modeling horizon but higher in later periods than would be calculated using the DICE2007 damage function. Specifically, the percent difference between damages in the base run of DICE2010 and those that would be calculated using the DICE2007 damage function. Specifically, the percent in 2005, decreases to +20 percent by 2300 (the end of the interagency analysis time horizon), and to +160 percent by the end of the model time horizon in 2595. The large increases in the far future years of the time horizon are due to the permanence associated with damages from sea level rise, along with the assumption that the sea level is projected to continue to rise long after the global average temperature begins to decrease. The changes to the loss function generally decrease the interagency working group SCC estimates slightly given that relative increases in later periods are discounted more heavily, all else equal.

B. FUND

FUND version 3.8 includes a number of changes over the previous version 3.5 (Narita et al. 2010) used in the 2010 interagency report. Documentation supporting FUND and the model's source code for all versions of the model is available from the model authors.⁸ Notable changes, due to their impact on the

⁷ The model and documentation, including formulas, are available on the author's webpage at <u>http://www.econ.yale.edu/~nordhaus/homepage/RICEmodels.htm</u>.

⁸ <u>http://www.fund-model.org/</u>. This report uses version 3.8 of the FUND model, which represents a modest update to the most recent version of the model to appear in the literature (version 3.7) (Anthoff and Tol, 2013a). For the purpose of computing the SCC, the relevant changes (between 3.7 to 3.8) are associated with improving

SCC estimates, are adjustments to the space heating, agriculture, and sea level rise damage functions in addition to changes to the temperature response function and the inclusion of indirect effects from methane emissions.⁹ We discuss each of these in turn.

Space Heating

In FUND, the damages associated with the change in energy needs for space heating are based on the estimated impact due to one degree of warming. These baseline damages are scaled based on the forecasted temperature anomaly's deviation from the one degree benchmark and adjusted for changes in vulnerability due to economic and energy efficiency growth. In FUND 3.5, the function that scales the base year damages adjusted for vulnerability allows for the possibility that in some simulations the benefits associated with reduced heating needs may be an unbounded convex function of the temperature anomaly. In FUND 3.8, the form of the scaling has been modified to ensure that the function is everywhere concave and that there will exist an upper bound on the benefits a region may receive from reduced space heating needs. The new formulation approaches a value of two in the limit of large temperature anomalies, or in other words, assuming no decrease in vulnerability, the reduced expenditures on space heating at any level of warming will not exceed two times the reductions experienced at one degree of warming. Since the reduced need for space heating represents a benefit of climate change in the model, or a negative damage, this change will increase the estimated SCC. This update accounts for a significant portion of the difference in the expected SCC estimates reported by the two versions of the model when run probabilistically.

Sea Level Rise and Land Loss

The FUND model explicitly includes damages associated with the inundation of dry land due to sea level rise. The amount of land lost within a region is dependent upon the proportion of the coastline being protected by adequate sea walls and the amount of sea level rise. In FUND 3.5 the function defining the potential land lost in a given year due to sea level rise is linear in the rate of sea level rise for that year. This assumption implicitly assumes that all regions are well represented by a homogeneous coastline in length and a constant uniform slope moving inland. In FUND 3.8 the function defining the potential land lost has been changed to be a convex function of sea level rise, thereby assuming that the slope of the shore line increases moving inland. The effect of this change is to typically reduce the vulnerability of some regions to sea level rise based land loss, thereby lowering the expected SCC estimate.¹⁰

Agriculture

consistency with IPCC AR4 by adjusting the atmospheric lifetimes of CH4 and N2O and incorporating the indirect forcing effects of CH4, along with making minor stability improvements in the sea wall construction algorithm. ⁹ The other damage sectors (water resources, space cooling, land loss, migration, ecosystems, human health, and

extreme weather) were not significantly updated.

¹⁰ For stability purposes this report also uses an update to the model which assumes that regional coastal protection measures will be built to protect the most valuable land first, such that the marginal benefits of coastal protection is decreasing in the level of protection following Fankhauser (1995).

In FUND, the damages associated with the agricultural sector are measured as proportional to the sector's value. The fraction is bounded from above by one and is made up of three additive components that represent the effects from carbon fertilization, the rate of temperature change, and the level of the temperature anomaly. In both FUND 3.5 and FUND 3.8, the fraction of the sector's value lost due to the level of the temperature anomaly is modeled as a quadratic function with an intercept of zero. In FUND 3.5, the coefficients of this loss function are modeled as the ratio of two random normal variables. This specification had the potential for unintended extreme behavior as draws from the parameter in the denominator approached zero or went negative. In FUND 3.8, the coefficients are drawn directly from truncated normal distributions so that they remain in the range $[0,\infty)$ and $(-\infty,0]$, respectively, ensuring the correct sign and eliminating the potential for divide by zero errors. The means for the new distributions are set equal to the ratio of the means from the normal distributions used in the previous version. In general the impact of this change has been to decrease the range of the distribution while spreading out the distributions' mass over the remaining range relative to the previous version. The net effect of this change on the SCC estimates is difficult to predict.

Transient Temperature Response

The temperature response model translates changes in global levels of radiative forcing into the current expected temperature anomaly. In FUND, a given year's increase in the temperature anomaly is based on a mean reverting function where the mean equals the equilibrium temperature anomaly that would eventually be reached if that year's level of radiative forcing were sustained. The rate of mean reversion defines the rate at which the transient temperature approaches the equilibrium. In FUND 3.5, the rate of temperature response is defined as a decreasing linear function of equilibrium climate sensitivity to capture the fact that the progressive heat uptake of the deep ocean causes the rate to slow at higher values of the equilibrium climate sensitivity. In FUND 3.8, the rate of temperature response has been updated to a quadratic function of the equilibrium climate sensitivity. This change reduces the sensitivity of the rate of temperature response to the level of the equilibrium climate sensitivity, a relationship first noted by Hansen et al. (1985) based on the heat uptake of the deep ocean. Therefore in FUND 3.8, the temperature response will typically be faster than in the previous version. The overall effect of this change is likely to increase estimates of the SCC as higher temperatures are reached during the timeframe analyzed and as the same damages experienced in the previous version of the model are now experienced earlier and therefore discounted less.

Methane

The IPCC AR4 notes a series of indirect effects of methane emissions, and has developed methods for proxying such effects when computing the global warming potential of methane (Forster et al. 2007). FUND 3.8 now includes the same methods for incorporating the indirect effects of methane emissions. Specifically, the average atmospheric lifetime of methane has been set to 12 years to account for the feedback of methane emissions on its own lifetime. The radiative forcing associated with atmospheric methane has also been increased by 40% to account for its net impact on ozone production and stratospheric water vapor. All else equal, the effect of this increased radiative forcing will be to increase the estimated SCC values, due to greater projected temperature anomaly.

C. PAGE

PAGE09 (Hope 2013) includes a number of changes from PAGE2002, the version used in the 2010 SCC interagency report. The changes that most directly affect the SCC estimates include: explicitly modeling the impacts from sea level rise, revisions to the damage function to ensure damages are constrained by GDP, a change in the regional scaling of damages, a revised treatment for the probability of a discontinuity within the damage function, and revised assumptions on adaptation. The model also includes revisions to the carbon cycle feedback and the calculation of regional temperatures.¹¹ More details on PAGE09 can be found in Hope (2011a, 2011b, 2011c). A description of PAGE2002 can be found in Hope (2006).

Sea Level Rise

While PAGE2002 aggregates all damages into two categories – economic and non-economic impacts -, PAGE09 adds a third explicit category: damages from sea level rise. In the previous version of the model, damages from sea level rise were subsumed by the other damage categories. In PAGE09 sea level damages increase less than linearly with sea level under the assumption that land, people, and GDP are more concentrated in low-lying shoreline areas. Damages from the economic and non-economic sector were adjusted to account for the introduction of this new category.

Revised Damage Function to Account for Saturation

In PAGE09, small initial economic and non-economic benefits (negative damages) are modeled for small temperature increases, but all regions eventually experience economic damages from climate change, where damages are the sum of additively separable polynomial functions of temperature and sea level rise. Damages transition from this polynomial function to a logistic path once they exceed a certain proportion of remaining Gross Domestic Product (GDP) to ensure that damages do not exceed 100 percent of GDP. This differs from PAGE2002, which allowed Eastern Europe to potentially experience large benefits from temperature increases, and which also did not bound the possible damages that could be experienced.

Regional Scaling Factors

As in the previous version of PAGE, the PAGE09 model calculates the damages for the European Union (EU) and then, assumes that damages for other regions are proportional based on a given scaling factor. The scaling factor in PAGE09 is based on the length of a region's coastline relative to the EU (Hope 2011b). Because of the long coastline in the EU, other regions are, on average, less vulnerable than the EU for the same sea level and temperature increase, but all regions have a positive scaling factor. PAGE2002 based its scaling factors on four studies reported in the IPCC's third assessment report, and allowed for benefits from temperature increase in Eastern Europe, smaller impacts in developed countries, and higher damages in developing countries.

¹¹ Because several changes in the PAGE model are structural (e.g., the addition of sea level rise and treatment of discontinuity), it is not possible to assess the direct impact of each change on the SCC in isolation as done for the other two models above.

Probability of a Discontinuity

In PAGE2002, the damages associated with a "discontinuity" (nonlinear extreme event) were modeled as an expected value. Specifically, a stochastic probability of a discontinuity was multiplied by the damages associated with a discontinuity to obtain an expected value, and this was added to the economic and non-economic impacts. That is, additional damages from an extreme event, such as extreme melting of the Greenland ice sheet, were multiplied by the probability of the event occurring and added to the damage estimate. In PAGE09, the probability of discontinuity is treated as a discrete event for each year in the model. The damages for each model run are estimated either with or without a discontinuity occurring, rather than as an expected value. A large-scale discontinuity becomes possible when the temperature rises beyond some threshold value between 2 and 4°C. The probability that a discontinuity will occur beyond this threshold then increases by between 10 and 30 percent for every 1°C rise in temperature beyond the threshold. If a discontinuity occurs, the EU loses an additional 5 to 25 percent of its GDP (drawn from a triangular distribution with a mean of 15 percent) in addition to other damages, and other regions lose an amount determined by the regional scaling factor. The threshold value for a possible discontinuity is lower than in PAGE2002, while the rate at which the probability of a discontinuity increases with the temperature anomaly and the damages that result from a discontinuity are both higher than in PAGE2002. The model assumes that only one discontinuity can occur and that the impact is phased in over a period of time, but once it occurs, its effect is permanent.

Adaptation

As in PAGE2002, adaptation is available to help mitigate any climate change impacts that occur. In PAGE this adaptation is the same regardless of the temperature change or sea level rise and is therefore akin to what is more commonly considered a reduction in vulnerability. It is modeled by reducing the damages by some percentage. PAGE09 assumes a smaller decrease in vulnerability than the previous version of the model and assumes that it will take longer for this change in vulnerability to be realized. In the aggregated economic sector, at the time of full implementation, this adaptation will mitigate all damages up to a temperature increase of 1°C, and for temperature anomalies between 1°C and 2°C, it will reduce damages by 15-30 percent (depending on the region). However, it takes 20 years to fully implement this adaptation. In PAGE2002, adaptation was assumed to reduce economic sector damages up to 2°C by 50-90 percent after 20 years. Beyond 2°C, no adaptation is assumed to be available to mitigate the impacts of climate change. For the non-economic sector, in PAGE09 adaptation is available to reduce 15 percent of the damages due to a temperature increase between 0°C and 2°C and is assumed to take 40 years to fully implement, instead of 25 percent of the damages over 20 years assumed in PAGE2002. Similarly, adaptation is assumed to alleviate 25-50 percent of the damages from the first 0.20 to 0.25 meters of sea level rise but is assumed to be ineffective thereafter. Hope (2011c) estimates that the less optimistic assumptions regarding the ability to offset impacts of temperature and sea level rise via adaptation increase the SCC by approximately 30 percent.

Other Noteworthy Changes

Two other changes in the model are worth noting. There is a change in the way the model accounts for decreased CO₂ absorption on land and in the ocean as temperature rises. PAGE09 introduces a linear feedback from global mean temperature to the percentage gain in the excess concentration of CO₂, capped at a maximum level. In PAGE2002, an additional amount was added to the CO₂ emissions each period to account for a decrease in ocean absorption and a loss of soil carbon. Also updated is the method by which the average global and annual temperature anomaly is downscaled to determine annual average regional temperature anomalies to be used in the regional damage functions. In PAGE2002, the scaling was determined solely based on regional difference in emissions of sulfate aerosols. In PAGE09, this regional temperature anomaly is further adjusted using an additive factor that is based on the average absolute latitude of a region relative to the area weighted average absolute latitude of the Earth's landmass, to capture relatively greater changes in temperature forecast to be experienced at higher latitudes.

III. Revised SCC Estimates

The updated versions of the three integrated assessment models were run using the same methodology detailed in the 2010 TSD (Interagency Working Group on Social Cost of Carbon 2010). The approach along with the inputs for the socioeconomic emissions scenarios, equilibrium climate sensitivity distribution, and discount rate remains the same. This includes the five reference scenarios based on the EMF-22 modeling exercise, the Roe and Baker equilibrium climate sensitivity distribution calibrated to the IPCC AR4, and three constant discount rates of 2.5, 3, and 5 percent.

As was previously the case, the use of three models, three discount rates, and five scenarios produces 45 separate distributions for the global SCC. The approach laid out in the 2010 TSD applied equal weight to each model and socioeconomic scenario in order to reduce the dimensionality down to three separate distributions representative of the three discount rates. The interagency group selected four values from these distributions for use in regulatory analysis. Three values are based on the average SCC across models and socio-economic-emissions scenarios at the 2.5, 3, and 5 percent discount rates, respectively. The fourth value was chosen to represent the higher-than-expected economic impacts from climate change further out in the tails of the SCC distribution. For this purpose, the 95th percentile of the SCC estimates at a 3 percent discount rate was chosen. (A detailed set of percentiles by model and scenario combination and additional summary statistics for the 2020 values is available in the Appendix.) As noted in the 2010 TSD, "the 3 percent discount rate is the central value, and so the central value that emerges is the average SCC across models at the 3 percent discount rate" (Interagency Working Group on Social Cost of Carbon 2010, p. 25). However, for purposes of capturing the uncertainties involved in regulatory impact analysis, the interagency group emphasizes the importance and value of including all four SCC values.

Table 2 shows the four selected SCC estimates in five year increments from 2010 to 2050. Values for 2010, 2020, 2030, 2040, and 2050 are calculated by first combining all outputs (10,000 estimates per


model run) from all scenarios and models for a given discount rate. Values for the years in between are calculated using linear interpolation. The full set of revised annual SCC estimates between 2010 and 2050 is reported in the Appendix.

			•	
Discount Rate	5.0%	3.0%	2.5%	3.0%
Year	Avg	Avg	Avg	95th
2010	11	32	51	89
2015	11	37	57	109
2020	12	43	64	128
2025	14	47	69	143
2030	16	52	75	159
2035	19	56	80	175
2040	21	61	86	191
2045	24	66	92	206
2050	26	71	97	220

Table 2: Revised Social Cost of CO₂, 2010 – 2050 (in 2007 dollars per metric ton of CO₂)

The SCC estimates using the updated versions of the models are higher than those reported in the 2010 TSD due to the changes to the models outlined in the previous section. By way of comparison, the 2020 SCC estimates reported in the original TSD were \$7, \$26, \$42 and \$81 (2007\$) (Interagency Working Group on Social Cost of Carbon 2010). Figure 1 illustrates where the four SCC values for 2020 fall within the full distribution for each discount rate based on the combined set of runs for each model and scenario (150,000 estimates in total for each discount rate). In general, the distributions are skewed to the right and have long tails. The Figure also shows that the lower the discount rate, the longer the right tail of the distribution.

Figure 1: Distribution of SCC Estimates for 2020 (in 2007\$ per metric ton CO₂)



As was the case in the 2010 TSD, the SCC increases over time because future emissions are expected to produce larger incremental damages as physical and economic systems become more stressed in response to greater climatic change. The approach taken by the interagency group is to compute the cost of a marginal ton emitted in the future by running the models for a set of perturbation years out to 2050. Table 3 illustrates how the growth rate for these four SCC estimates varies over time.

Average Annual Growth	5.0%	3.0%	2.5%	3.0%
Rate (%)	Avg	Avg	Avg	95th
2010-2020	1.2%	3.3%	2.4%	4.4%
2020-2030	3.4%	2.1%	1.7%	2.4%
2030-2040	3.0%	1.9%	1.5%	2.1%
2040-2050	2.6%	1.6%	1.3%	1.5%

Table 3: Average Annual Growth Rates of SCC Estimates between 2010 and 2050

The future monetized value of emission reductions in each year (the SCC in year t multiplied by the change in emissions in year t) must be discounted to the present to determine its total net present value for use in regulatory analysis. As previously discussed in the 2010 TSD, damages from future emissions should be discounted at the same rate as that used to calculate the SCC estimates themselves to ensure internal consistency – i.e., future damages from climate change, whether they result from emissions today or emissions in a later year, should be discounted using the same rate.

Under current OMB guidance contained in Circular A-4, analysis of economically significant proposed and final regulations from the domestic perspective is required, while analysis from the international perspective is optional. However, the climate change problem is highly unusual in at least two respects. First, it involves a global externality: emissions of most greenhouse gases contribute to damages around the world even when they are emitted in the United States. Consequently, to address the global nature of the problem, the SCC must incorporate the full (global) damages caused by GHG emissions. Second, climate change presents a problem that the United States alone cannot solve. Even if the United States were to reduce its greenhouse gas emissions to zero, that step would be far from enough to avoid substantial climate change. Other countries would also need to take action to reduce emissions if significant changes in the global climate are to be avoided. Emphasizing the need for a global solution to a global problem, the United States has been actively involved in seeking international agreements to reduce emissions and in encouraging other nations, including emerging major economies, to take significant steps to reduce emissions. When these considerations are taken as a whole, the interagency group concluded that a global measure of the benefits from reducing U.S. emissions is preferable. For additional discussion, see the 2010 TSD.

IV. Other Model Limitations and Research Gaps

The 2010 interagency SCC TSD discusses a number of important limitations for which additional research is needed. In particular, the document highlights the need to improve the quantification of both noncatastrophic and catastrophic damages, the treatment of adaptation and technological change, and the way in which inter-regional and inter-sectoral linkages are modeled. While the new version of the models discussed above offer some improvements in these areas, further work remains warranted. The 2010 TSD also discusses the need to more carefully assess the implications of risk aversion for SCC estimation as well as the inability to perfectly substitute between climate and non-climate goods at higher temperature increases, both of which have implications for the discount rate used. EPA, DOE, and other agencies continue to engage in research on modeling and valuation of climate impacts that can potentially improve SCC estimation in the future.

References

Anthoff, D. and Tol, R.S.J. 2013a. The uncertainty about the social cost of carbon: a decomposition analysis using FUND. *Climatic Change* 117: 515–530.

Anthoff, D. and Tol, R.S.J. 2013b. Erratum to: The uncertainty about the social cost of carbon: A decomposition analysis using FUND. *Climatic Change*. Advance online publication. doi: 10.1007/s10584-013-0959-1.

Fankhauser, S. 1995. Valuing climate change: The economics of the greenhouse. London, England: Earthscan.

Forster, P., V. Ramaswamy, P. Artaxo, T. Berntsen, R. Betts, D.W. Fahey, J. Haywood, J. Lean, D.C. Lowe, G. Myhre, J. Nganga, R. Prinn, G. Raga, M. Schulz and R. Van Dorland. 2007. Changes in Atmospheric Constituents and in Radiative Forcing. In: *Climate Change 2007: The Physical Science Basis. Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change* [Solomon, S., D. Qin, M. Manning, Z. Chen, M. Marquis, K.B. Averyt, M.Tignor and H.L. Miller (eds.)]. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA.

Hope, Chris. 2006. "The Marginal Impact of CO_2 from PAGE2002: An Integrated Assessment Model Incorporating the IPCC's Five Reasons for Concern." *The Integrated Assessment Journal*. 6(1): 19–56.

Hope, Chris. 2011a "The PAGE09 Integrated Assessment Model: A Technical Description" Cambridge Judge Business School Working Paper No. 4/2011 (April). Accessed November 23, 2011: http://www.jbs.cam.ac.uk/research/working papers/2011/wp1104.pdf.

Hope, Chris. 2011b "The Social Cost of CO₂ from the PAGE09 Model" Cambridge Judge Business School Working Paper No. 5/2011 (June). Accessed November 23, 2011: <u>http://www.jbs.cam.ac.uk/research/working_papers/2011/wp1105.pdf</u>.

Hope, Chris. 2011c "New Insights from the PAGE09 Model: The Social Cost of CO₂" Cambridge Judge Business School Working Paper No. 8/2011 (July). Accessed November 23, 2011: <u>http://www.jbs.cam.ac.uk/research/working papers/2011/wp1108.pdf</u>.

Hope, C. 2013. Critical issues for the calculation of the social cost of CO₂: why the estimates from PAGE09 are higher than those from PAGE2002. *Climatic Change* 117: 531–543.

Interagency Working Group on Social Cost of Carbon. 2010. Social Cost of Carbon for Regulatory Impact Analysis under Executive Order 12866. February. United States Government. http://www.whitehouse.gov/sites/default/files/omb/inforeg/for-agencies/Social-Cost-of-Carbon-for-RIA.pdf.

Meehl, G.A., T.F. Stocker, W.D. Collins, P. Friedlingstein, A.T. Gaye, J.M. Gregory, A. Kitoh, R. Knutti, J.M. Murphy, A. Noda, S.C.B. Raper, I.G. Watterson, A.J. Weaver and Z.-C. Zhao. 2007. Global Climate Projections. In: *Climate Change 2007: The Physical Science Basis. Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change* [Solomon, S., D. Qin, M. Manning, Z. Chen, M. Marquis, K.B. Averyt, M. Tignor and H.L. Miller (eds.)]. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA.

Narita, D., R. S. J. Tol and D. Anthoff. 2010. Economic costs of extratropical storms under climate change: an application of FUND. *Journal of Environmental Planning and Management* 53(3): 371-384.

National Academy of Sciences. 2011. *Climate Stabilization Targets: Emissions, Concentrations, and Impacts over Decades to Millennia*. Washington, DC: National Academies Press, Inc.

Nicholls, R.J., N. Marinova, J.A. Lowe, S. Brown, P. Vellinga, D. de Gusmão, J. Hinkel and R.S.J. Tol. 2011. Sea-level rise and its possible impacts given a 'beyond 4°C world' in the twenty-first century. *Phil. Trans. R. Soc. A* 369(1934): 161-181.

Nordhaus, W. 2010. Economic aspects of global warming in a post-Copenhagen environment. *Proceedings of the National Academy of Sciences* 107(26): 11721-11726.

Nordhaus, W. 2008. A Question of Balance: Weighing the Options on Global Warming Policies. New Haven, CT: Yale University Press.

Randall, D.A., R.A. Wood, S. Bony, R. Colman, T. Fichefet, J. Fyfe, V. Kattsov, A. Pitman, J. Shukla, J. Srinivasan, R.J. Stouffer, A. Sumi and K.E. Taylor. 2007. Climate Models and Their Evaluation. In: *Climate Change 2007: The Physical Science Basis. Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change* [Solomon, S., D. Qin, M. Manning, Z. Chen, M. Marquis, K.B. Averyt, M.Tignor and H.L. Miller (eds.)]. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA.

Appendix A

Discount Rate	5.0%	3.0%	2.5%	3.0%
Year	Avg	Avg	Avg	95th
2010	11	32	51	89
2011	11	33	52	93
2012	11	34	54	97
2013	11	35	55	101
2014	11	36	56	105
2015	11	37	57	109
2016	12	38	59	112
2017	12	39	60	116
2018	12	40	61	120
2019	12	42	62	124
2020	12	43	64	128
2021	12	43	65	131
2022	13	44	66	134
2023	13	45	67	137
2024	14	46	68	140
2025	14	47	69	143
2026	15	48	70	146
2027	15	49	71	149
2028	15	50	72	152
2029	16	51	73	155
2030	16	52	75	159
2031	17	52	76	162
2032	17	53	77	165
2033	18	54	78	168
2034	18	55	79	172
2035	19	56	80	175
2036	19	57	81	178
2037	20	58	83	181
2038	20	59	84	185
2039	21	60	85	188
2040	21	61	86	191
2041	22	62	87	194
2042	22	63	88	197
2043	23	64	89	200
2044	23	65	90	203
2045	24	66	92	206
2046	24	67	93	209
2047	25	68	94	211
2048	25	69	95	214
2049	26	70	96	217
2050	26	71	97	220

Table A1: Annual SCC Values: 2010-2050 (2007\$/metric ton CO₂)

Percentile	1st	5th	10th	25th	50th	Avg	75th	90th	95 th	99th
Scenario ¹²					PA	GE				
IMAGE	6	11	15	27	58	129	139	327	515	991
MERGE	4	6	9	16	34	78	82	196	317	649
MESSAGE	4	8	11	20	42	108	107	278	483	918
MiniCAM Base	5	9	12	22	47	107	113	266	431	872
5th Scenario	2	4	6	11	25	85	68	200	387	955
		-	-	-						
Scenario					DI	CE				
IMAGE	25	31	37	47	64	72	92	123	139	161
MERGE	14	18	20	26	36	40	50	65	74	85
MESSAGE	20	24	28	37	51	58	71	95	109	221
MiniCAM Base	20	25	29	38	53	61	76	102	117	135
5th Scenario	17	22	25	33	45	52	65	91	106	126
Scenario		-	-	-	FU	ND	-			
IMAGE	-14	-2	4	15	31	39	55	86	107	157
MERGE	-6	1	6	14	27	35	46	70	87	141
MESSAGE	-16	-5	1	11	24	31	43	67	83	126
MiniCAM Base	-7	2	7	16	32	39	55	83	103	158
5th Scenario	-29	-13	-6	4	16	21	32	53	69	103

Table A2: 2020 Global SCC Estimates at 2.5 Percent Discount Rate (2007\$/metric ton CO₂)

Table A3: 2020 Global SCC Estimates at 3 Percent Discount Rate (2007\$/metric ton CO₂)

Percentile	1st	5th	10th	25th	50th	Avg	75th	90th	95th	99th
Scenario	PAGE									
IMAGE	4	7	10	18	38	91	95	238	385	727
MERGE	2	4	6	11	23	56	58	142	232	481
MESSAGE	3	5	7	13	29	75	74	197	330	641
MiniCAM Base	3	5	8	14	30	73	75	184	300	623
5th Scenario	1	3	4	7	17	58	48	136	264	660
r	-	-	-			-	-	-		
Scenario					DI	CE				
IMAGE	16	21	24	32	43	48	60	79	90	102
MERGE	10	13	15	19	25	28	35	44	50	58
MESSAGE	14	18	20	26	35	40	49	64	73	83
MiniCAM Base	13	17	20	26	35	39	49	65	73	85
5th Scenario	12	15	17	22	30	34	43	58	67	79
		-						-	-	
Scenario		-	-		FU	ND	-	-		-
IMAGE	-13	-4	0	8	18	23	33	51	65	99
MERGE	-7	-1	2	8	17	21	29	45	57	95
MESSAGE	-14	-6	-2	5	14	18	26	41	52	82
MiniCAM Base	-7	-1	3	9	19	23	33	50	63	101
5th Scenario	-22	-11	-6	1	8	11	18	31	40	62

¹² See 2010 TSD for a description of these scenarios.

Dorcontilo	1.0+	Гtb	10+h	JE+b	F O+h	Ava	7C+b	00+h	0 C +b	00+h
Percentile	1st	5th	<u>10th</u>	25th	<u>50th</u>	Avg	75th	90th	95th	99th
Scenario										
IMAGE	1	2	2	5	10	28	27	71	123	244
MERGE	1	1	2	3	7	17	17	45	75	153
MESSAGE	1	1	2	4	9	24	22	60	106	216
MiniCAM Base	1	1	2	3	8	21	21	54	94	190
5th Scenario	0	1	1	2	5	18	14	41	78	208
			-	-			-	-		
Scenario					DI	CE				
IMAGE	6	8	9	11	14	15	18	22	25	27
MERGE	4	5	6	7	9	10	12	15	16	18
MESSAGE	6	7	8	10	12	13	16	20	22	25
MiniCAM Base	5	6	7	8	11	12	14	18	20	22
5th Scenario	5	6	6	8	10	11	14	17	19	21
Scenario		-	-		FU	ND	-	-		
IMAGE	-9	-5	-4	-1	2	3	6	10	14	24
MERGE	-6	-4	-2	0	3	4	6	11	15	26
MESSAGE	-10	-6	-4	-1	1	2	5	9	12	21
MiniCAM Base	-7	-4	-2	0	3	4	6	11	14	25
5th Scenario	-11	-7	-5	-3	0	0	3	5	7	13

Table A4: 2020 Global SCC Estimates at 5 Percent Discount Rate (2007\$/metric ton CO₂)

Discount rate:	count rate: 5.0%					3.0%				2.5%			
Statistic:	Mean	Variance	Skewness	Kurtosis	Mean	Variance	Skewness	Kurtosis	Mean	Variance	Skewness	Kurtosis	
DICE	12	26	2	15	38	409	3	24	57	1097	3	30	
PAGE	22	1616	5	32	71	14953	4	22	101	29312	4	23	
FUND	3	41	5	179	19	1452	-42	8727	33	6154	-73	14931	

Table A5: Additional Summary Statistics of 2020 Global SCC Estimates

Appendix B

The November 2013 revision of this technical support document is based on two corrections to the runs based on the FUND model. First, the potential dry land loss in the algorithm that estimates regional coastal protections was misspecified in the model's computer code. This correction is covered in an erratum to Anthoff and Tol (2013a) published in the same journal (Climatic Change) in October 2013 (Anthoff and Tol (2013b)). Second, the equilibrium climate sensitivity distribution was inadvertently specified as a truncated Gamma distribution (the default in FUND) as opposed to the truncated Roe and Baker distribution as was intended. The truncated Gamma distribution used in the FUND runs had approximately the same mean and upper truncation point, but lower variance and faster decay of the upper tail, as compared to the intended specification based on the Roe and Baker distribution. The difference between the original estimates reported in the May 2013 version of this technical support document and this revision are generally one dollar or less.