

**Technical Support Document: -
Technical Update of the Social Cost of Carbon for Regulatory Impact Analysis -
Under Executive Order 12866 -**

Interagency Working Group on Social Cost of Greenhouse Gases, United States Government

With participation by

Council of Economic Advisers
Council on Environmental Quality
Department of Agriculture
Department of Commerce
Department of Energy
Department of the Interior
Department of Transportation
Department of the Treasury
Environmental Protection Agency
National Economic Council
Office of Management and Budget
Office of Science and Technology Policy

August 2016

See Appendix B for Details on Revisions since May 2013

Preface

The Interagency Working Group on the Social Cost of Greenhouse Gases (formerly the Interagency Working Group on the Social Cost of Carbon) has a longstanding commitment to ensure that the social cost of carbon estimates continue to reflect the best available science and methodologies. Given this commitment and public comments on issues of a deeply technical nature received by the Office of Management and Budget and federal agencies, the Interagency Working Group is seeking independent expert advice on technical opportunities to update the social cost of carbon estimates. The Interagency Working Group asked the National Academies of Sciences, Engineering, and Medicine in 2015 to review the latest research on modeling the economic aspects of climate change to inform future revisions to the social cost of carbon estimates presented in this technical support document. In January 2016, the Academies' Committee on the Social Cost of Carbon issued an interim report that recommended against a near-term update to the social cost of carbon estimates, but included recommendations for enhancing the presentation and discussion of uncertainty around the current estimates. This revision to the TSD responds to these recommendations in the presentation of the current estimates. It does not revisit the interagency group's 2010 methodological decisions or update the schedule of social cost of carbon estimates presented in the July 2015 revision. The Academies' final report (expected in early 2017) will provide longer term recommendations for a more comprehensive update.

Executive Summary

Executive Order 12866 requires agencies, to the extent permitted by law, “to assess both the costs and the benefits of the intended regulation and, recognizing that some costs and benefits are difficult to quantify, propose or adopt a regulation only upon a reasoned determination that the benefits of the intended regulation justify its costs.” The purpose of the social cost of carbon (SC-CO₂)¹ estimates presented here is to allow agencies to incorporate the social benefits of reducing carbon dioxide (CO₂) emissions into cost-benefit analyses of regulatory actions. The SC-CO₂ is the monetized damages associated with an incremental increase in carbon emissions in a given year. It is intended to include (but is not limited to) changes in net agricultural productivity, human health, property damages from increased flood risk, and the value of ecosystem services due to climate change.

The interagency process that developed the original U.S. government SC-CO₂ estimates is described in the 2010 Technical Support Document on the Social Cost of Carbon (TSD) (Interagency Working Group on Social Cost of Carbon 2010). Through that process the Interagency Working Group (IWG) selected SC-CO₂ values for use in regulatory analyses. For each emissions year, four values are recommended. Three of these values are based on the average SC-CO₂ from three integrated assessment models (IAMs), at discount rates of 2.5, 3, and 5 percent. In addition, as discussed in the 2010 TSD, there is extensive evidence in the scientific and economic literature on the potential for lower-probability, but higher-impact outcomes from climate change, which would be particularly harmful to society and thus relevant to the public and policymakers. The fourth value is thus included to represent the marginal damages associated with these lower-probability, higher-impact outcomes. Accordingly, this fourth value is selected from further out in the tail of the distribution of SC-CO₂ estimates; specifically, the fourth value corresponds to the 95th percentile of the frequency distribution of SC-CO₂ estimates based on a 3 percent discount rate. Because the present value of economic damages associated with CO₂ emissions change over time, a separate set of estimates is presented for each emissions year through 2050, which is sufficient to cover the time frame addressed in most current regulatory impact analyses.

In May of 2013, the IWG provided an update of the SC-CO₂ estimates based on new versions of each IAM (DICE, PAGE, and FUND). The 2013 update did not revisit other IWG modeling decisions (e.g., the discount rate, reference case socioeconomic and emission scenarios, or equilibrium climate sensitivity). Improvements in the way damages are modeled were confined to those that had been incorporated into the latest versions of the models by the developers themselves in the peer-reviewed literature. The IWG subsequently provided additional minor technical revisions in November of 2013 and July of 2015, as described in Appendix B.

The purpose of this 2016 revision to the TSD is to enhance the presentation and discussion of quantified uncertainty around the current SC-CO₂ estimates, as a response to recommendations in the interim report by the National Academies of Sciences, Engineering, and Medicine. Included herein are an expanded

¹ Throughout this Technical Support Document (TSD) we refer to the estimates as “SC-CO₂ estimates” rather than the more simplified “SCC” abbreviation used in previous versions of the TSD.

graphical presentation of the SC-CO₂ estimates highlighting a symmetric range of uncertainty around estimates for each discount rate, new sections that provide a unified discussion of the methodology used to incorporate sources of uncertainty, and a detailed explanation of the uncertain parameters in both the FUND and PAGE models.

The distributions of SC-CO₂ estimates reflect uncertainty in key model parameters chosen by the IWG such as the sensitivity of the climate to increases in carbon dioxide concentrations, as well as uncertainty in default parameters set by the original model developers. This TSD maintains the same approach to estimating the SC-CO₂ and selecting four values for each emissions year that was used in earlier versions of the TSD. Table ES-1 summarizes the SC-CO₂ estimates for the years 2010 through 2050. These estimates are identical to those reported in the previous version of the TSD, released in July 2015. As explained in previous TSDs, the central value is the average of SC-CO₂ estimates based on the 3 percent discount rate. For purposes of capturing uncertainty around the SC-CO₂ estimates in regulatory impact analysis, the IWG emphasizes the importance of considering all four SC-CO₂ values.


Table ES-1: Social Cost of CO₂, 2010 – 2050 (in 2007 dollars per metric ton of CO₂)

Year	5% Average	3% Average	2.5% Average	High Impact (95 th Pct at 3%)
2010	10	31	50	86
2015	11	36	56	105
2020	12	42	62	123
2025	14	46	68	138
2030	16	50	73	152
2035	18	55	78	168
2040	21	60	84	183
2045	23	64	89	197
2050	26	69	95	212

While point estimates are important for providing analysts with a tractable approach for regulatory analysis, they do not fully quantify uncertainty associated with the SC-CO₂ estimates. Figure ES-1 presents the quantified sources of uncertainty in the form of frequency distributions for the SC-CO₂ estimates for emissions in 2020. To highlight the difference between the impact of the discount rate on the SC-CO₂ and other quantified sources of uncertainty, the bars below the frequency distributions provide a symmetric representation of quantified variability in the SC-CO₂ estimates for each discount rate. When an agency determines that it is appropriate to conduct additional quantitative uncertainty analysis, it should follow best practices for probabilistic analysis.² The full set of information that underlies the frequency distributions in Figure ES-1, which have previously been available upon request, are now available on Office of Management and Budget’s (OMB) website for easy public access.

² See e.g. OMB Circular A-4, section on *Treatment of Uncertainty*. Available at: https://www.whitehouse.gov/omb/circulars_a004_a-4/#e.

Figure ES-1: Frequency Distribution of SC-CO₂ Estimates for 2020³



³ Although the distributions in Figure ES-1 are based on the full set of model results (150,000 estimates for each discount rate), for display purposes the horizontal axis is truncated with 0.1 to 0.6 percent of the estimates lying below the lowest bin displayed and 0.2 to 3.7 percent of the estimates lying above the highest bin displayed, depending on the discount rate.

I. Purpose

The purpose of this document is to present the current schedule of social cost of carbon (SC-CO₂) estimates, along with an enhanced presentation and discussion of quantified sources of uncertainty around the estimates to respond to recommendations in the interim report of the National Academies of Sciences, Engineering, and Medicine (National Academies 2016).⁴ Because the last substantive update to the SC-CO₂ estimates occurred in May 2013, this document maintains much of the earlier technical discussion from the May 2013 TSD. The SC-CO₂ estimates themselves remain unchanged since the July 2015 revision.

E.O. 13563 commits the Administration to regulatory decision making “based on the best available science.”⁵ Additionally, the IWG recommended in 2010 that the SC-CO₂ estimates be revisited on a regular basis or as model updates that reflect the growing body of scientific and economic knowledge become available.⁶ By early 2013, new versions of the three integrated assessment models (IAMs) used by the U.S. government to estimate the SC-CO₂ (DICE, FUND, and PAGE) were available and had been published in the peer-reviewed literature. While acknowledging the continued limitations of the approach taken by the IWG in 2010 (documented in the original 2010 TSD), the May 2013 TSD provided an update of the SC-CO₂ estimates based on the latest peer-reviewed version of the models, replacing model versions that were developed up to ten years earlier in a rapidly evolving field. It did not revisit other assumptions with regard to the discount rate, reference case socioeconomic and emission scenarios, or equilibrium climate sensitivity. Improvements in the way damages are modeled were confined to those that had been incorporated into the latest versions of the models by the developers themselves in the peer-reviewed literature. The agencies participating in the IWG continue to investigate potential improvements to the way in which economic damages associated with changes in CO₂ emissions are quantified.

Section II summarizes the major features of the IAMs used in this TSD that were updated in 2013 relative to the versions of the models used in the 2010 TSD. Section III presents the SC-CO₂ estimates for 2010 – 2050 based on these versions of the models. Section IV discusses the treatment of uncertainty in the analysis. Section V provides a discussion of other model limitations and research gaps.

II. Summary of Model Updates

This section briefly reviews the features of the three IAMs used in this TSD (DICE 2010, FUND 3.8, and PAGE 2009) that were updated by the model developers relative to the versions of the models used by the IWG in 2010 (DICE 2007, FUND 3.5, and PAGE 2002). The focus here is on describing those model updates that are relevant to estimating the social cost of carbon, as summarized in Table 1. For example, both the DICE and PAGE models now include an explicit representation of sea level rise damages. Other

⁴ In this document, we present all social cost estimates per metric ton of CO₂ emissions. Alternatively, one could report the social cost per metric ton of carbon emissions. The multiplier for translating between mass of CO₂ and the mass of carbon is 3.67 (the molecular weight of CO₂ divided by the molecular weight of carbon = 44/12 = 3.67).

⁵ http://www.whitehouse.gov/sites/default/files/omb/inforeg/eo12866/eo13563_01182011.pdf

⁶ See p. 1, 3, 4, 29, and 33 (Interagency Working Group on Social Cost of Carbon 2010).

revisions to PAGE include: updated adaptation assumptions, revisions to ensure damages are constrained by GDP, updated regional scaling of damages, and a revised treatment of potentially abrupt shifts in climate damages. The DICE model’s simple carbon cycle has been updated to be more consistent with a more complex climate model. The FUND model includes updated damage functions for sea level rise impacts, the agricultural sector, and reduced space heating requirements, as well as changes to the transient response of temperature to the buildup of GHG concentrations and the inclusion of indirect effects of methane emissions. Changes made to parts of the models that are superseded by the IWG’s modeling assumptions—regarding equilibrium climate sensitivity, discounting, and socioeconomic variables—are not discussed here but can be found in the references provided in each section below.

Table 1: Summary of Key Model Revisions Relevant to the IWG SC-CO₂ Estimates

IAM	Version used in 2010 IWG Analysis	Version Used since May 2013	Key changes relevant to IWG SC-CO₂
DICE	2007	2010	Updated calibration of the carbon cycle model and explicit representation of sea level rise (SLR) and associated damages.
FUND	3.5 (2009)	3.8 (2012)	Updated damage functions for space heating, SLR, agricultural impacts, changes to transient response of temperature to buildup of GHG concentrations, and inclusion of indirect climate effects of methane.
PAGE	2002	2009	Explicit representation of SLR damages, revisions to damage function to ensure damages do not exceed 100% of GDP, change in regional scaling of damages, revised treatment of potential abrupt damages, and updated adaptation assumptions.

A. DICE

DICE 2010 includes a number of changes over the previous 2007 version used in the 2010 TSD. The model changes that are relevant for the SC-CO₂ estimates developed by the IWG include: 1) updated parameter values for the carbon cycle model, 2) an explicit representation of sea level dynamics, and 3) a re-calibrated damage function that includes an explicit representation of economic damages from sea level rise. Changes were also made to other parts of the DICE model—including the equilibrium climate sensitivity parameter, the rate of change of total factor productivity, and the elasticity of the marginal utility of consumption—but these components of DICE are superseded by the IWG’s assumptions and so will not be discussed here. More details on DICE2007 can be found in Nordhaus (2008) and on DICE2010 in Nordhaus (2010). The DICE2010 model and documentation is also available for download from the homepage of William Nordhaus.

Carbon Cycle Parameters

DICE uses a three-box model of carbon stocks and flows to represent the accumulation and transfer of carbon among the atmosphere, the shallow ocean and terrestrial biosphere, and the deep ocean. These

parameters are “calibrated to match the carbon cycle in the Model for the Assessment of Greenhouse Gas Induced Climate Change (MAGICC)” (Nordhaus 2008, p. 44).⁷ Carbon cycle transfer coefficient values in DICE2010 are based on re-calibration of the model to match the newer 2009 version of MAGICC (Nordhaus 2010, p. 2). For example, in DICE2010, in each decade 12 percent of the carbon in the atmosphere is transferred to the shallow ocean, 4.7 percent of the carbon in the shallow ocean is transferred to the atmosphere, 94.8 percent remains in the shallow ocean, and 0.5 percent is transferred to the deep ocean. For comparison, in DICE 2007, 18.9 percent of the carbon in the atmosphere is transferred to the shallow ocean each decade, 9.7 percent of the carbon in the shallow ocean is transferred to the atmosphere, 85.3 percent remains in the shallow ocean, and 5 percent is transferred to the deep ocean.

The implication of these changes for DICE2010 is in general a weakening of the ocean as a carbon sink and therefore a higher concentration of carbon in the atmosphere than in DICE2007 for a given path of emissions. All else equal, these changes will generally increase the level of warming and therefore the SC-CO₂ estimates in DICE2010 relative to those from DICE2007.

Sea Level Dynamics

A new feature of DICE2010 is an explicit representation of the dynamics of the global average sea level anomaly to be used in the updated damage function (discussed below). This section contains a brief description of the sea level rise (SLR) module; a more detailed description can be found on the model developer’s website.⁸ The average global sea level anomaly is modeled as the sum of four terms that represent contributions from: 1) thermal expansion of the oceans, 2) melting of glaciers and small ice caps, 3) melting of the Greenland ice sheet, and 4) melting of the Antarctic ice sheet.

The parameters of the four components of the SLR module are calibrated to match consensus results from the IPCC’s Fourth Assessment Report (AR4).⁹ The rise in sea level from thermal expansion in each time period (decade) is 2 percent of the difference between the sea level in the previous period and the long run equilibrium sea level, which is 0.5 meters per degree Celsius (°C) above the average global temperature in 1900. The rise in sea level from the melting of glaciers and small ice caps occurs at a rate of 0.008 meters per decade per °C above the average global temperature in 1900.

The contribution to sea level rise from melting of the Greenland ice sheet is more complex. The equilibrium contribution to SLR is 0 meters for temperature anomalies less than 1 °C and increases linearly from 0 meters to a maximum of 7.3 meters for temperature anomalies between 1 °C and 3.5 °C. The contribution to SLR in each period is proportional to the difference between the previous period’s sea

⁷ MAGICC is a simple climate model initially developed by the U.S. National Center for Atmospheric Research that has been used heavily by the Intergovernmental Panel on Climate Change (IPCC) to emulate projections from more sophisticated state of the art earth system simulation models (Randall et al. 2007).

⁸ Documentation on the new sea level rise module of DICE is available on William Nordhaus’ website at: http://nordhaus.econ.yale.edu/documents/SLR_021910.pdf.

⁹ For a review of post-IPCC AR4 research on sea level rise, see Nicholls et al. (2011) and NAS (2011).

level anomaly and the equilibrium sea level anomaly, where the constant of proportionality increases with the temperature anomaly in the current period.

The contribution to SLR from the melting of the Antarctic ice sheet is -0.001 meters per decade when the temperature anomaly is below 3 °C and increases linearly between 3 °C and 6 °C to a maximum rate of 0.025 meters per decade at a temperature anomaly of 6 °C.

Re-calibrated Damage Function

Economic damages from climate change in the DICE model are represented by a fractional loss of gross economic output in each period. A portion of the remaining economic output in each period (net of climate change damages) is consumed and the remainder is invested in the physical capital stock to support future economic production, so each period's climate damages will reduce consumption in that period and in all future periods due to the lost investment. The fraction of output in each period that is lost due to climate change impacts is represented as a sigmoid, or "S"-shaped, function of the temperature anomaly in the period.¹⁰ The loss function in DICE2010 has been expanded by including a quadratic sub-function of SLR. In DICE2010 the temperature anomaly coefficients have been recalibrated to avoid double-counting damages from sea level rise that were implicitly included in these parameters in DICE2007.

The aggregate damages in DICE2010 are illustrated by Nordhaus (2010, p. 3), who notes that "...damages in the uncontrolled (baseline) [i.e., reference] case ... in 2095 are \$12 trillion, or 2.8 percent of global output, for a global temperature increase of 3.4 °C above 1900 levels." This compares to a loss of 3.2 percent of global output at 3.4 °C in DICE2007. However, in DICE2010, annual damages are lower in most of the early periods of the modeling horizon but higher in later periods than would be calculated using the DICE2007 damage function. Specifically, the percent difference between damages in the base run of DICE2010 and those that would be calculated using the DICE2007 damage function starts at +7 percent in 2005, decreases to a low of -14 percent in 2065, then continuously increases to +20 percent by 2300 (the end of the IWG analysis time horizon), and to +160 percent by the end of the model time horizon in 2595. The large increases in the far future years of the time horizon are due to the permanence associated with damages from sea level rise, along with the assumption that the sea level is projected to continue to rise long after the global average temperature begins to decrease. The changes to the loss function generally decrease the IWG SC-CO₂ estimates slightly given that relative increases in damages in later periods are discounted more heavily, all else equal.

B. FUND

FUND version 3.8 includes a number of changes over the previous version 3.5 (Narita et al. 2010) used in the 2010 TSD. Documentation supporting FUND and the model's source code for all versions of the model

¹⁰ The model and documentation, including formulas, are available on the author's webpage at <http://www.econ.yale.edu/~nordhaus/homepage/RICEmodels.htm>.

is available from the model authors.¹¹ Notable changes, due to their impact on the SC-CO₂ estimates, are adjustments to the space heating, agriculture, and sea level rise damage functions in addition to changes to the temperature response function and the inclusion of indirect effects from methane emissions.¹² Each of these is discussed in turn.

Space Heating

In FUND, the damages associated with the change in energy needs for space heating are based on the estimated impact due to one degree of warming. These baseline damages are scaled based on the forecasted temperature anomaly's deviation from the one degree benchmark and adjusted for changes in vulnerability due to economic and energy efficiency growth. In FUND 3.5, the function that scales the base year damages adjusted for vulnerability allows for the possibility that in some simulations the benefits associated with reduced heating needs may be an unbounded convex function of the temperature anomaly. In FUND 3.8, the form of the scaling has been modified to ensure that the function is everywhere concave and that there will exist an upper bound on the benefits a region may receive from reduced space heating needs. The new formulation approaches a value of two in the limit of large temperature anomalies, or in other words, assuming no decrease in vulnerability, the reduced expenditures on space heating at any level of warming will not exceed two times the reductions experienced at one degree of warming. Since the reduced need for space heating represents a benefit of climate change in the model, or a negative damage, this change will increase the estimated SC-CO₂. This update accounts for a significant portion of the difference in the expected SC-CO₂ estimates reported by the two versions of the model when run probabilistically.

Sea Level Rise and Land Loss

The FUND model explicitly includes damages associated with the inundation of dry land due to sea level rise. The amount of land lost within a region depends on the proportion of the coastline being protected by adequate sea walls and the amount of sea level rise. In FUND 3.5 the function defining the potential land lost in a given year due to sea level rise is linear in the rate of sea level rise for that year. This assumption implicitly assumes that all regions are well represented by a homogeneous coastline in length and a constant uniform slope moving inland. In FUND 3.8 the function defining the potential land lost has been changed to be a convex function of sea level rise, thereby assuming that the slope of the shore line

¹¹ <http://www.fund-model.org/>. This report uses version 3.8 of the FUND model, which represents a modest update to the most recent version of the model to appear in the literature (version 3.7) (Anthoff and Tol, 2013a, 2013b). For the purpose of computing the SC-CO₂, the relevant changes (between 3.7 to 3.8) are associated with improving consistency with IPCC AR4 by adjusting the atmospheric lifetimes of CH₄ and N₂O and incorporating the indirect forcing effects of CH₄, along with making minor stability improvements in the sea wall construction algorithm.

¹² The other damage sectors (water resources, space cooling, land loss, migration, ecosystems, human health, and extreme weather) were not significantly updated.

increases moving inland. The effect of this change is to typically reduce the vulnerability of some regions to sea level rise based land loss, thereby lowering the expected SC-CO₂ estimate.¹³

¹³ For stability purposes this report also uses an update to the model which assumes that regional coastal protection measures will be built to protect the most valuable land first, such that the marginal benefits of coastal protection is decreasing in the level of protection following Fankhauser (1995).

Agriculture

In FUND, the damages associated with the agricultural sector are measured as proportional to the sector's value. The fraction is bounded from above by one and is made up of three additive components that represent the effects from carbon fertilization, the rate of temperature change, and the level of the temperature anomaly. In both FUND 3.5 and FUND 3.8, the fraction of the sector's value lost due to the level of the temperature anomaly is modeled as a quadratic function with an intercept of zero. In FUND 3.5, the coefficients of this loss function are modeled as the ratio of two random normal variables. This specification had the potential for unintended extreme behavior as draws from the parameter in the denominator approached zero or went negative. In FUND 3.8, the coefficients are drawn directly from truncated normal distributions so that they remain in the range $[0, \infty)$ and $(-\infty, 0]$, respectively, ensuring the correct sign and eliminating the potential for divide-by-zero errors. The means for the new distributions are set equal to the ratio of the means from the normal distributions used in the previous version. In general the impact of this change has been to decrease the range of the distribution while spreading out the distributions' mass over the remaining range relative to the previous version. The net effect of this change on the SC-CO₂ estimates is difficult to predict.

Transient Temperature Response

The temperature response model translates changes in global levels of radiative forcing into the current expected temperature anomaly. In FUND, a given year's increase in the temperature anomaly is based on a mean reverting function where the mean equals the equilibrium temperature anomaly that would eventually be reached if that year's level of radiative forcing were sustained. The rate of mean reversion defines the rate at which the transient temperature approaches the equilibrium. In FUND 3.5, the rate of temperature response is defined as a decreasing linear function of equilibrium climate sensitivity to capture the fact that the progressive heat uptake of the deep ocean causes the rate to slow at higher values of the equilibrium climate sensitivity. In FUND 3.8, the rate of temperature response has been updated to a quadratic function of the equilibrium climate sensitivity. This change reduces the sensitivity of the rate of temperature response to the level of the equilibrium climate sensitivity, a relationship first noted by Hansen et al. (1985) based on the heat uptake of the deep ocean. Therefore in FUND 3.8, the temperature response will typically be faster than in the previous version. The overall effect of this change is likely to increase estimates of the SC-CO₂ as higher temperatures are reached during the timeframe analyzed and as the same damages experienced in the previous version of the model are now experienced earlier and therefore discounted less.

Methane

The IPCC AR4 notes a series of indirect effects of methane emissions, and has developed methods for proxying such effects when computing the global warming potential of methane (Forster et al. 2007). FUND 3.8 now includes the same methods for incorporating the indirect effects of methane emissions. Specifically, the average atmospheric lifetime of methane has been set to 12 years to account for the feedback of methane emissions on its own lifetime. The radiative forcing associated with atmospheric methane has also been increased by 40% to account for its net impact on ozone production and

stratospheric water vapor. This update to the model is relevant for the SC-CO₂ because most of the damage functions are non-linear functions of the temperature anomaly, which represents the fact that as the climate system becomes more stressed an additional unit of warming will have a greater impact on damages. Accounting for the indirect effects of CH₄ emissions on temperature will therefore move the model further up the damage curves in the baseline, making a marginal change in emissions of CO₂ more impactful. All else equal, the effect of this increased radiative forcing will be to increase the estimated SC-CO₂ values, due to greater projected temperature anomaly.

C. PAGE

PAGE09 (Hope 2013) includes a number of changes from PAGE2002, the version used in the 2010 TSD. The changes that most directly affect the SC-CO₂ estimates include: explicitly modeling the impacts from sea level rise, revisions to the damage function to ensure damages are constrained by GDP, a change in the regional scaling of damages, a revised treatment for the probability of a discontinuity within the damage function, and revised assumptions on adaptation. The model also includes revisions to the carbon cycle feedback and the calculation of regional temperatures.¹⁴ More details on PAGE09 can be found in Hope (2011a, 2011b, 2011c). A description of PAGE2002 can be found in Hope (2006).

Sea Level Rise

While PAGE2002 aggregates all damages into two categories—economic and non-economic impacts—PAGE09 adds a third explicit category: damages from sea level rise. In the previous version of the model, damages from sea level rise were subsumed by the other damage categories. In PAGE09 sea level damages increase less than linearly with sea level under the assumption that land, people, and GDP are more concentrated in low-lying shoreline areas. Damages from the economic and non-economic sectors were adjusted to account for the introduction of this new category.

Revised Damage Function to Account for Saturation

In PAGE09, small initial economic and non-economic benefits (negative damages) are modeled for small temperature increases, but all regions eventually experience economic damages from climate change, where damages are the sum of additively separable polynomial functions of temperature and sea level rise. Damages transition from this polynomial function to a logistic path once they exceed a certain proportion of remaining Gross Domestic Product (GDP) to ensure that damages do not exceed 100 percent of GDP. This differs from PAGE2002, which allowed Eastern Europe to potentially experience large benefits from temperature increases, and which also did not bound the possible damages that could be experienced.

¹⁴ Because several changes in the PAGE model are structural (e.g., the addition of sea level rise and treatment of discontinuity), it is not possible to assess the direct impact of each change on the SC-CO₂ in isolation as done for the other two models above.

Regional Scaling Factors

As in the previous version of PAGE, the PAGE09 model calculates the damages for the European Union (EU) and then, assumes that damages for other regions are proportional based on a given scaling factor. The scaling factors in PAGE09 are based on the length of each region's coastline relative to the EU (Hope 2011b). Because of the long coastline in the EU, other regions are, on average, less vulnerable than the EU for the same sea level and temperature increase, but all regions have a positive scaling factor. PAGE2002 based its scaling factors on four studies reported in the IPCC's third assessment report, and allowed for benefits from temperature increases in Eastern Europe, smaller impacts in developed countries, and higher damages in developing countries.

Probability of a Discontinuity

In PAGE2002, the damages associated with a "discontinuity" (nonlinear extreme event) were modeled as an expected value. Specifically, a stochastic probability of a discontinuity was multiplied by the damages associated with a discontinuity to obtain an expected value, and this was added to the economic and non-economic impacts. That is, additional damages from an extreme event, such as extreme melting of the Greenland ice sheet, were multiplied by the probability of the event occurring and added to the damage estimate. In PAGE09, the probability of discontinuity is treated as a discrete event for each year in the model. The damages for each model run are estimated either with or without a discontinuity occurring, rather than as an expected value. A large-scale discontinuity becomes possible when the temperature rises beyond some threshold value between 2 and 4°C. The probability that a discontinuity will occur beyond this threshold then increases by between 10 and 30 percent for every 1°C rise in temperature beyond the threshold. If a discontinuity occurs, the EU loses an additional 5 to 25 percent of its GDP (drawn from a triangular distribution with a mean of 15 percent) in addition to other damages, and other regions lose an amount determined by their regional scaling factor. The threshold value for a possible discontinuity is lower than in PAGE2002, while the rate at which the probability of a discontinuity increases with the temperature anomaly and the damages that result from a discontinuity are both higher than in PAGE2002. The model assumes that only one discontinuity can occur and that the impact is phased in over a period of time, but once it occurs, its effect is permanent.

Adaptation

As in PAGE2002, adaptation is available to help mitigate any climate change impacts that occur. In PAGE this adaptation is the same regardless of the temperature change or sea level rise and is therefore akin to what is more commonly considered a reduction in vulnerability. It is modeled by reducing the damages by some percentage. PAGE09 assumes a smaller decrease in vulnerability than the previous version of the model and assumes that it will take longer for this change in vulnerability to be realized. In the aggregated economic sector, at the time of full implementation, this adaptation will mitigate all damages up to a temperature increase of 1°C, and for temperature anomalies between 1°C and 2°C, it will reduce damages by 15-30 percent (depending on the region). However, it takes 20 years to fully implement this adaptation. In PAGE2002, adaptation was assumed to reduce economic sector damages up to 2°C by 50-90 percent after 20 years. Beyond 2°C, no adaptation is assumed to be available to mitigate the impacts of climate

change. For the non-economic sector, in PAGE09 adaptation is available to reduce 15 percent of the damages due to a temperature increase between 0°C and 2°C and is assumed to take 40 years to fully implement, instead of 25 percent of the damages over 20 years assumed in PAGE2002. Similarly, adaptation is assumed to alleviate 25-50 percent of the damages from the first 0.20 to 0.25 meters of sea level rise but is assumed to be ineffective thereafter. Hope (2011c) estimates that the less optimistic assumptions regarding the ability to offset impacts of temperature and sea level rise via adaptation increase the SC-CO₂ by approximately 30 percent.

Other Noteworthy Changes

Two other changes in the model are worth noting. There is a change in the way the model accounts for decreased CO₂ absorption on land and in the ocean as temperature rises. PAGE09 introduces a linear feedback from global mean temperature to the percentage gain in the excess concentration of CO₂, capped at a maximum level. In PAGE2002, an additional amount was added to the CO₂ emissions each period to account for a decrease in ocean absorption and a loss of soil carbon. Also updated is the method by which the average global and annual temperature anomaly is downscaled to determine annual average regional temperature anomalies to be used in the regional damage functions. In PAGE2002, the scaling was determined solely based on regional difference in emissions of sulfate aerosols. In PAGE09, this regional temperature anomaly is further adjusted using an additive factor that is based on the average absolute latitude of a region relative to the area weighted average absolute latitude of the Earth's landmass, to capture relatively greater changes in temperature forecast to be experienced at higher latitudes.

III. SC-CO₂ Estimates

The three IAMs were run using the same methodology detailed in the 2010 TSD (Interagency Working Group on Social Cost of Carbon 2010). The approach, along with the inputs for the socioeconomic emissions scenarios, equilibrium climate sensitivity distribution, and discount rate remains the same. This includes the five reference scenarios based on the EMF-22 modeling exercise, the Roe and Baker equilibrium climate sensitivity distribution calibrated to the IPCC AR4, and three constant discount rates of 2.5, 3, and 5 percent.

As was previously the case, use of three models, three discount rates, and five scenarios produces 45 separate frequency distributions of SC-CO₂ estimates in a given year. The approach laid out in the 2010 TSD applied equal weight to each model and socioeconomic scenario in order to reduce the dimensionality down to three separate distributions, one for each of the three discount rates. The IWG selected four values from these distributions for use in regulatory analysis. Three values are based on the average SC-CO₂ across models and socioeconomic and emissions scenarios at the 2.5, 3, and 5 percent discount rates, respectively. The fourth value is included to provide information on the marginal damages associated with lower-probability, higher-impact outcomes that would be particularly harmful to society. As discussed in the 2010 TSD, there is extensive evidence in the scientific and economic literature of the potential for lower-probability, higher-impact outcomes from climate change, which would be particularly harmful to society and thus relevant to the public and policymakers. This points to the relevance of values above the

mean in right skewed distributions. Accordingly, this fourth value is selected from further out in the tails of the frequency distribution of SC-CO₂ estimates, and, in particular, is set to the 95th percentile of the frequency distribution of SC-CO₂ estimates based on a 3 percent discount rate. (A detailed set of percentiles by model and scenario combination and additional summary statistics for the 2020 values is available in Appendix A.) As noted in the 2010 TSD, “the 3 percent discount rate is the central value, and so the central value that emerges is the average SC-CO₂ across models at the 3 percent discount rate” (Interagency Working Group on Social Cost of Carbon 2010, p. 25). However, for purposes of capturing the uncertainties involved in regulatory impact analysis, the IWG emphasizes the importance and value of including all four SC-CO₂ values.

Table 2 shows the four selected SC-CO₂ estimates in five year increments from 2010 to 2050. Values for 2010, 2020, 2030, 2040, and 2050 are calculated by first combining all outputs (10,000 estimates per model run) from all scenarios and models for a given discount rate. Values for the years in between are calculated using linear interpolation. The full set of revised annual SC-CO₂ estimates between 2010 and 2050 is reported in the Appendix and the full set of model results are available on the OMB website.¹⁵

Table 2: Social Cost of CO₂, 2010 – 2050 (in 2007 dollars per metric ton of CO₂)

Year	5% Average	3% Average	2.5% Average	High Impact (95 th Pct at 3%)
2010	10	31	50	86
2015	11	36	56	105
2020	12	42	62	123
2025	14	46	68	138
2030	16	50	73	152
2035	18	55	78	168
2040	21	60	84	183
2045	23	64	89	197
2050	26	69	95	212

As was the case in the 2010 TSD, the SC-CO₂ increases over time because future emissions are expected to produce larger incremental damages as physical and economic systems become more stressed in response to greater climatic change, and because GDP is growing over time and many damage categories are modeled as proportional to gross GDP. The approach taken by the IWG is to compute the cost of a marginal ton emitted in the future by running the models for a set of perturbation years out to 2050. Table 3 illustrates how the growth rate for these four SC-CO₂ estimates varies over time.

¹⁵ <https://www.whitehouse.gov/omb/oira/social-cost-of-carbon>.

Table 3: Average Annual Growth Rates of SC-CO₂ Estimates between 2010 and 2050

Average Annual Growth Rate (%)	5.0% Avg	3.0% Avg	2.5% Avg	3.0% 95th
2010-2020	1.2%	3.2%	2.4%	4.4%
2020-2030	3.4%	2.1%	1.7%	2.3%
2030-2040	3.0%	1.9%	1.5%	2.0%
2040-2050	2.6%	1.6%	1.3%	1.6%

The future monetized value of emission reductions in each year (the SC-CO₂ in year *t* multiplied by the change in emissions in year *t*) must be discounted to the present to determine its total net present value for use in regulatory analysis. As previously discussed in the 2010 TSD, damages from future emissions should be discounted at the same rate as that used to calculate the SC-CO₂ estimates themselves to ensure internal consistency—i.e., future damages from climate change, whether they result from emissions today or emissions in a later year, should be discounted to the base year of the analysis using the same rate.

Current guidance contained in OMB Circular A-4 indicates that analysis of economically significant proposed and final regulations from the domestic perspective is required, while analysis from the international perspective is optional. However, the IWG (including OMB) determined that a modified approach is more appropriate in this case because the climate change problem is highly unusual in a number of respects. First, it involves a global externality: emissions of most greenhouse gases contribute to damages around the world even when they are emitted in the United States—and conversely, greenhouse gases emitted elsewhere contribute to damages in the United States. Consequently, to address the global nature of the problem, the SC-CO₂ must incorporate the full (global) damages caused by GHG emissions. Second, climate change presents a problem that the United States alone cannot solve. Other countries will also need to take action to reduce emissions if significant changes in the global climate are to be avoided. Emphasizing the need for a global solution to a global problem, the United States has been actively involved in seeking international agreements to reduce emissions. For example, the United States joined over 170 other nations and signed the Paris Agreement on April 22, 2016, signaling worldwide commitment to reduce GHG emissions. The United States has been active in encouraging other nations, including emerging major economies, to take significant steps to reduce emissions. Using a global estimate of damages in U.S. regulatory analyses sends a strong signal to other nations that they too should base their emissions reductions strategies on a global perspective, thus supporting a cooperative and mutually beneficial approach to achieving needed reduction. Thirteen prominent academics noted that these "are compelling reasons to focus on a global [SC-CO₂]" in a recent article on the SC-CO₂ (Pizer et al. 2014). In addition, adverse impacts on other countries can have spillover effects on the United States, particularly in the areas of national security, international trade, public health, and humanitarian concerns. When these considerations are taken as a whole, the IWG concluded that a global measure of the benefits from reducing U.S. emissions is appropriate. For additional discussion, see the 2010 TSD.

IV. Treatment of Uncertainty

Uncertainty about the value of the SC-CO₂ is in part inherent, as with any analysis that looks into the future, but it is also driven by current data gaps associated with the complex physical, economic, and behavioral processes that link GHG emissions to human health and well-being. Some sources of uncertainty pertain to aspects of the natural world, such as quantifying the physical effects of greenhouse gas emissions on Earth systems. Other sources of uncertainty are associated with current and future human behavior and well-being, such as population and economic growth, GHG emissions, the translation of Earth system changes to economic damages, and the role of adaptation. It is important to note that even in the presence of uncertainty, scientific and economic analysis can provide valuable information to the public and decision makers, though the uncertainty should be acknowledged and when possible taken into account in the analysis. This section summarizes the sources of uncertainty that the IWG was able to consider in a quantitative manner in estimating the SC-CO₂. Further discussion on sources of uncertainty that are active areas of research and have not yet been fully quantified in the SC-CO₂ estimates is provided in Section V and in the 2010 TSD.

In developing the SC-CO₂ estimates, the IWG considered various sources of uncertainty through a combination of a multi-model ensemble, probabilistic analysis, and scenario analysis. For example, the three IAMs used collectively span a wide range of Earth system and economic outcomes to help reflect the uncertainty in the literature and in the underlying dynamics being modeled. The use of an ensemble of three different models is also intended to, at least partially, address the fact that no single model includes all of the quantified economic damages. It also helps to reflect structural uncertainty across the models, which is uncertainty in the underlying relationships between GHG emissions, Earth systems, and economic damages that are included in the models. Bearing in mind the different limitations of each model (discussed in the 2010 TSD) and lacking an objective basis upon which to differentially weight the models, the three IAMs are given equal weight in the analysis.

The IWG used Monte Carlo techniques to run the IAMs a large number of times. In each simulation the uncertain parameters are represented by random draws from their defined probability distributions. In all three models the equilibrium climate sensitivity is treated probabilistically based on the probability distribution described in the 2010 TSD. The equilibrium climate sensitivity is a key parameter in this analysis because it helps define the strength of the climate response to increasing GHG concentrations in the atmosphere. In addition, the FUND and PAGE models define many of their parameters with probability distributions instead of point estimates. For these two models, the model developers' default probability distributions are maintained for all parameters other than those superseded by the IWG's harmonized inputs (i.e., equilibrium climate sensitivity, socioeconomic and emissions scenarios, and discount rates). More information on the uncertain parameters in PAGE and FUND is presented in Appendix C.

For the socioeconomic and emissions scenarios, uncertainty is included in the analysis by considering a range of scenarios, which are described in detail in the 2010 SC-CO₂ TSD. As noted in the 2010 TSD, while the IWG considered formally assigning probability weights to the different socioeconomic scenarios selected, it came to the conclusion that this could not be accomplished in an analytically rigorous way given the dearth of information on the likelihood of a full range of future socioeconomic pathways. Thus,

the IWG determined that, because no basis for assigning differential weights was available, the most transparent way to present a range of uncertainty was simply to weight each of the five scenarios equally for the consolidated estimates. To provide additional information as to how the results vary with the scenarios, summarized results for each scenario are presented separately in Appendix A. The results of each model run are available on the OMB website.


Finally, based on the review of the literature, the IWG chose discount rates that reflect reasonable judgements under both prescriptive and descriptive approaches to intergenerational discounting. As discussed in the 2010 TSD, in light of disagreement in the literature on the appropriate discount rate to use in this context and uncertainty about how rates may change over time, the IWG selected three certainty-equivalent constant discount rates to span a plausible range: 2.5, 3, and 5 percent per year. However, unlike the approach taken for consolidating results across models and socioeconomic and emissions scenarios, the SC-CO₂ estimates are not pooled across different discount rates because the range of discount rates reflects both uncertainty and, at least in part, different policy or value judgements.

The outcome of accounting for various sources of uncertainty using the approaches described above is a frequency distribution of the SC-CO₂ estimates for emissions occurring in a given year for each of the three discount rates. These frequency distributions reflect the uncertainty around the input parameters for which probability distributions were defined, as well as from the multi-model ensemble and socioeconomic and emissions scenarios where probabilities were implied by the equal weighting assumption. It is important to note that the set of SC-CO₂ estimates obtained from this analysis does not yield a probability distribution that fully characterizes uncertainty about the SC-CO₂ due to impact categories omitted from the models and sources of uncertainty that have not been fully characterized due to data limitations.

Figure 1 presents the frequency distribution of the SC-CO₂ estimates for emissions in 2020 for each of the three discount rates. Each of these distributions represents 150,000 estimates based on 10,000 simulations for each combination of the three models and five socioeconomic and emissions scenarios.¹⁶ In general, the distributions are skewed to the right and have long right tails, which tend to be even longer for lower discount rates. To highlight the difference between the impact of the discount rate on the SC-CO₂ and other quantified sources of uncertainty, the bars below the frequency distributions provide a symmetric representation of quantified variability in the SC-CO₂ estimates conditioned on each discount rate. The full set of SC-CO₂ results through 2050 is available on OMB's website. This may be useful to analysts in situations that warrant additional quantitative uncertainty analysis (e.g., as recommended by OMB for rules that exceed \$1 billion in annual benefits or costs). See OMB Circular A-4 for guidance and discussion of best practices in conducting uncertainty analysis in RIAs.

¹⁶ Although the distributions in Figure 1 are based on the full set of model results (150,000 estimates for each discount rate), for display purposes the horizontal axis is truncated with 0.1 to 0.6 percent of the estimates lying below the lowest bin displayed and 0.2 to 3.7 percent of the estimates lying above the highest bin displayed, depending on the discount rate.

Figure 1: Frequency Distribution of SC-CO₂ Estimates for 2020 (in 2007\$ per metric ton CO₂)



As previously described, the SC-CO₂ estimates produced by the IWG are based on a rigorous approach to accounting for quantifiable uncertainty using multiple analytical techniques. In addition, the scientific and economics literature has further explored known sources of uncertainty related to estimates of the SC-CO₂. For example, researchers have published papers that explore the sensitivity of IAMs and the resulting SC-CO₂ estimates to different assumptions embedded in the models (see, e.g., Hope (2013), Anthoff and Tol (2013a), and Nordhaus (2014)). However, there remain additional sources of uncertainty that have not been fully characterized and explored due to remaining data limitations. Additional research is needed in order to expand the quantification of various sources of uncertainty in estimates of the SC-CO₂ (e.g., developing explicit probability distributions for more inputs pertaining to climate impacts and their valuation). The IWG is actively following advances in the scientific and economic literature that could provide guidance on, or methodologies for, a more robust incorporation of uncertainty.

V. Other Model Limitations and Research Gaps

The 2010 SC-CO₂ TSD discusses a number of important limitations for which additional research is needed. In particular, the document highlights the need to improve the quantification of both non-catastrophic and catastrophic damages, the treatment of adaptation and technological change, and the way in which inter-regional and inter-sectoral linkages are modeled. While the more recent versions of the models discussed above offer some improvements in these areas, further research is still needed. Currently, IAMs do not include all of the important physical, ecological, and economic impacts of climate change

recognized in the climate change literature due to a lack of precise information on the nature of damages and because the science incorporated into these models understandably lags behind the most recent research.¹⁷ These individual limitations do not all work in the same direction in terms of their influence on the SC-CO₂ estimates; however, it is the IWG's judgment that, taken together, these limitations suggest that the SC-CO₂ estimates are likely conservative. In particular, the IPCC Fourth Assessment Report (Meehl et al. 2007), which was the most current IPCC assessment available at the time of the IWG's 2009-2010 review, concluded that SC-CO₂ estimates "very likely...underestimate the damage costs" due to omitted impacts. Since then, the peer-reviewed literature has continued to support this conclusion, as noted in the IPCC Fifth Assessment report (Oppenheimer et al. 2014).

Another area of active research relates to intergenerational discounting, including the application of discount rates to regulations in which some costs and benefits accrue intra-generationally while others accrue inter-generationally. Some experts have argued that a declining discount rate would be appropriate to analyze impacts that occur far into the future (Arrow et al. 2013). However, additional research and analysis is still needed to develop a methodology for implementing a declining discount rate and to understand the implications of applying these theoretical lessons in practice.

The 2010 TSD also discusses the need to more carefully assess the implications of risk aversion for SC-CO₂ estimation as well as the substitution possibilities between climate and non-climate goods at higher temperature increases, both of which have implications for the discount rate used. EPA, DOE, and other agencies continue to engage in research on modeling and valuation of climate impacts that can potentially improve SC-CO₂ estimation in the future. See the 2010 SC-CO₂ TSD for the full discussion.

¹⁷ See, for example, Howard (2014) and EPRI (2014) for recent discussions.

References

- Anthoff, D. and Tol, R.S.J. 2013a. The uncertainty about the social cost of carbon: a decomposition analysis using FUND. *Climatic Change* 117: 515–530.
- Anthoff, D. and Tol, R.S.J. 2013b. Erratum to: The uncertainty about the social cost of carbon: A decomposition analysis using FUND. *Climatic Change*. Advance online publication. doi: 10.1007/s10584-013-0959-1.
- Arrow, K., M. Cropper, C. Gollier, B. Groom, G. Heal, R. Newell, W. Nordhaus, R. Pindyck, W. Pizer, P. Portney, T. Sterner, R.S.J. Tol, and M. Weitzman. 2013. Determining Benefits and Costs for Future Generations. *Science* 341: 349-350.
- Electric Power Research Institute. 2014. Understanding the Social Cost of carbon: A Technical Assessment. www.epri.com.
- Fankhauser, S. 1995. Valuing climate change: The economics of the greenhouse. London, England: Earthscan.
- Forster, P., V. Ramaswamy, P. Artaxo, T. Berntsen, R. Betts, D.W. Fahey, J. Haywood, J. Lean, D.C. Lowe, G. Myhre, J. Nganga, R. Prinn, G. Raga, M. Schulz and R. Van Dorland. 2007. Changes in Atmospheric Constituents and in Radiative Forcing. In: *Climate Change 2007: The Physical Science Basis. Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change* [Solomon, S., D. Qin, M. Manning, Z. Chen, M. Marquis, K.B. Averyt, M. Tignor and H.L. Miller (eds.)]. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA.
- Hansen, J., G. Russell, A. Lacis, I. Fung, D. Rind, and P. Stone. 1985. Climate Response Times: Dependence on climate sensitivity and ocean mixing. *Science* 229: 857–859.
- Hope, Chris. 2006. “The Marginal Impact of CO₂ from PAGE2002: An Integrated Assessment Model Incorporating the IPCC’s Five Reasons for Concern.” *The Integrated Assessment Journal*. 6(1): 19–56.
- Hope, Chris. 2011a. “The PAGE09 Integrated Assessment Model: A Technical Description” Cambridge Judge Business School Working Paper No. 4/2011 (April). Accessed November 23, 2011: http://www.jbs.cam.ac.uk/research/working_papers/2011/wp1104.pdf.
- Hope, Chris. 2011b. “The Social Cost of CO₂ from the PAGE09 Model” Cambridge Judge Business School Working Paper No. 5/2011 (June). Accessed November 23, 2011: http://www.jbs.cam.ac.uk/research/working_papers/2011/wp1105.pdf.
- Hope, Chris. 2011c. “New Insights from the PAGE09 Model: The Social Cost of CO₂” Cambridge Judge Business School Working Paper No. 8/2011 (July). Accessed November 23, 2011: http://www.jbs.cam.ac.uk/research/working_papers/2011/wp1108.pdf.
- Hope, C. 2013. Critical issues for the calculation of the social cost of CO₂: why the estimates from PAGE09 are higher than those from PAGE2002. *Climatic Change* 117: 531–543.

Howard, Peter. 2014. Omitted Damages: What's Missing from the Social Cost of Carbon. [http://costofcarbon.org/files/Omitted Damages Whats Missing From the Social Cost of Carbon.pdf](http://costofcarbon.org/files/Omitted_Damages_Whats_Missing_From_the_Social_Cost_of_Carbon.pdf).

Interagency Working Group on Social Cost of Carbon. 2010. Technical Support Document: Social Cost of Carbon for Regulatory Impact Analysis under Executive Order 12866. February. United States Government. <http://www.whitehouse.gov/sites/default/files/omb/infoereg/for-agencies/Social-Cost-of-Carbon-for-RIA.pdf>.

Meehl, G.A., T.F. Stocker, W.D. Collins, P. Friedlingstein, A.T. Gaye, J.M. Gregory, A. Kitoh, R. Knutti, J.M. Murphy, A. Noda, S.C.B. Raper, I.G. Watterson, A.J. Weaver and Z.-C. Zhao. 2007. Global Climate Projections. In: *Climate Change 2007: The Physical Science Basis. Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change* [Solomon, S., D. Qin, M. Manning, Z. Chen, M. Marquis, K.B. Averyt, M. Tignor and H.L. Miller (eds.)]. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA.

Narita, D., R. S. J. Tol and D. Anthoff. 2010. Economic costs of extratropical storms under climate change: an application of FUND. *Journal of Environmental Planning and Management* 53(3): 371-384.

National Academy of Sciences (NAS). 2011. *Climate Stabilization Targets: Emissions, Concentrations, and Impacts over Decades to Millennia*. Washington, DC: National Academies Press, Inc.

National Academies of Sciences, Engineering, and Medicine (National Academies). 2016. Assessment of Approaches to Updating the Social Cost of Carbon: Phase 1 Report on a Near-Term Update. Washington, D.C.: National Academies Press. Available at: [http://sites.nationalacademies.org/DBASSE/BECS/Social Cost of Carbon Near Term Update/index.htm](http://sites.nationalacademies.org/DBASSE/BECS/Social_Cost_of_Carbon_Near_Term_Update/index.htm).

Nicholls, R.J., N. Marinova, J.A. Lowe, S. Brown, P. Vellinga, D. de Gusmão, J. Hinkel and R.S.J. Tol. 2011. Sea-level rise and its possible impacts given a 'beyond 4°C world' in the twenty-first century. *Phil. Trans. R. Soc. A* 369(1934): 161-181.

Nordhaus, W. 2014. Estimates of the Social Cost of Carbon: Concepts and Results from the DICE-2013R Model and Alternative Approaches. *Journal of the Association of Environmental and Resource Economists* 1(1/2): 273-312.

Nordhaus, W. 2010. Economic aspects of global warming in a post-Copenhagen environment. *Proceedings of the National Academy of Sciences* 107(26): 11721-11726.

Nordhaus, W. 2008. *A Question of Balance: Weighing the Options on Global Warming Policies*. New Haven, CT: Yale University Press.

Oppenheimer, M., M. Campos, R. Warren, J. Birkmann, G. Luber, B. O'Neill, and K. Takahashi, 2014: Emergent risks and key vulnerabilities. In: *Climate Change 2014: Impacts, Adaptation, and Vulnerability. Part A: Global and Sectoral Aspects. Contribution of Working Group II to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change* [Field, C.B., V.R. Barros, D.J. Dokken, K.J. Mach, M.D. Mastrandrea, T.E. Bilir, M. Chatterjee, K.L. Ebi, Y.O. Estrada, R.C. Genova, B. Girma, E.S. Kissel, A.N. Levy, S. MacCracken, P.R. Mastrandrea, and L.L. White (eds.)]. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, pp. 1039-1099.

Pizer, W., M. Adler, J. Aldy, D. Anthoff, M. Cropper, K. Gillingham, M. Greenstone, B. Murray, R. Newell, R. Richels, A. Rowell, S. Waldhoff, and J. Wiener. 2014. Using and Improving the Social Cost of Carbon. *Science* 346: 1181–82.

Randall, D.A., R.A. Wood, S. Bony, R. Colman, T. Fichet, J. Fyfe, V. Kattsov, A. Pitman, J. Shukla, J. Srinivasan, R.J. Stouffer, A. Sumi and K.E. Taylor. 2007. Climate Models and Their Evaluation. In: *Climate Change 2007: The Physical Science Basis. Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change* [Solomon, S., D. Qin, M. Manning, Z. Chen, M. Marquis, K.B. Averyt, M. Tignor and H.L. Miller (eds.)]. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA.

Appendix A

Table A1: Annual SC-CO₂ Values: 2010-2050 (2007\$/metric ton CO₂)

Year	5% Average	3% Average	2.5% Average	High Impact (95 th Pct at 3%)
2010	10	31	50	86
2011	11	32	51	90
2012	11	33	53	93
2013	11	34	54	97
2014	11	35	55	101
2015	11	36	56	105
2016	11	38	57	108
2017	11	39	59	112
2018	12	40	60	116
2019	12	41	61	120
2020	12	42	62	123
2021	12	42	63	126
2022	13	43	64	129
2023	13	44	65	132
2024	13	45	66	135
2025	14	46	68	138
2026	14	47	69	141
2027	15	48	70	143
2028	15	49	71	146
2029	15	49	72	149
2030	16	50	73	152
2031	16	51	74	155
2032	17	52	75	158
2033	17	53	76	161
2034	18	54	77	164
2035	18	55	78	168
2036	19	56	79	171
2037	19	57	81	174
2038	20	58	82	177
2039	20	59	83	180
2040	21	60	84	183
2041	21	61	85	186
2042	22	61	86	189
2043	22	62	87	192
2044	23	63	88	194
2045	23	64	89	197
2046	24	65	90	200
2047	24	66	92	203
2048	25	67	93	206
2049	25	68	94	209
2050	26	69	95	212

Table A2: 2020 Global SC-CO₂ Estimates at 2.5 Percent Discount Rate (2007\$/metric ton CO₂)

Percentile	1st	5th	10th	25th	50th	Avg	75th	90th	95th	99th
Scenario ¹⁸	PAGE									
IMAGE	6	10	15	26	55	123	133	313	493	949
MERGE Optimistic	4	6	8	15	32	75	79	188	304	621
MESSAGE	4	7	10	19	41	104	103	266	463	879
MiniCAM Base	5	8	12	21	45	102	108	255	412	835
5th Scenario	2	4	6	11	24	81	66	192	371	915

Scenario	DICE									
IMAGE	25	31	37	47	64	72	92	123	139	161
MERGE Optimistic	14	18	20	26	36	40	50	65	74	85
MESSAGE	20	24	28	37	51	58	71	95	109	221
MiniCAM Base	20	25	29	38	53	61	76	102	117	135
5th Scenario	17	22	25	33	45	52	65	91	106	126

Scenario	FUND									
IMAGE	-14	-2	4	15	31	39	55	86	107	157
MERGE Optimistic	-6	1	6	14	27	35	46	70	87	141
MESSAGE	-16	-5	1	11	24	31	43	67	83	126
MiniCAM Base	-7	2	7	16	32	39	55	83	103	158
5th Scenario	-29	-13	-6	4	16	21	32	53	69	103

Table A3: 2020 Global SC-CO₂ Estimates at 3 Percent Discount Rate (2007\$/metric ton CO₂)

Percentile	1st	5th	10th	25th	50th	Avg	75th	90th	95th	99th
Scenario	PAGE									
IMAGE	4	7	9	17	36	87	91	228	369	696
MERGE Optimistic	2	4	6	10	22	54	55	136	222	461
MESSAGE	3	5	7	13	28	72	71	188	316	614
MiniCAM Base	3	5	7	13	29	70	72	177	288	597
5th Scenario	1	3	4	7	16	55	46	130	252	632

Scenario	DICE									
IMAGE	16	21	24	32	43	48	60	79	90	102
MERGE Optimistic	10	13	15	19	25	28	35	44	50	58
MESSAGE	14	18	20	26	35	40	49	64	73	83
MiniCAM Base	13	17	20	26	35	39	49	65	73	85
5th Scenario	12	15	17	22	30	34	43	58	67	79

Scenario	FUND									
IMAGE	-13	-4	0	8	18	23	33	51	65	99
MERGE Optimistic	-7	-1	2	8	17	21	29	45	57	95
MESSAGE	-14	-6	-2	5	14	18	26	41	52	82
MiniCAM Base	-7	-1	3	9	19	23	33	50	63	101
5th Scenario	-22	-11	-6	1	8	11	18	31	40	62

¹⁸ See 2010 TSD for a description of these scenarios.

Table A4: 2020 Global SC-CO₂ Estimates at 5 Percent Discount Rate (2007\$/metric ton CO₂)

Percentile	1st	5th	10th	25th	50th	Avg	75th	90th	95th	99th
Scenario	PAGE									
IMAGE	1	2	2	4	10	27	26	68	118	234
MERGE Optimistic	1	1	2	3	6	17	17	43	72	146
MESSAGE	1	1	2	4	8	23	22	58	102	207
MiniCAM Base	1	1	2	3	8	20	20	52	90	182
5th Scenario	0	1	1	2	5	17	14	39	75	199

Scenario	DICE									
IMAGE	6	8	9	11	14	15	18	22	25	27
MERGE Optimistic	4	5	6	7	9	10	12	15	16	18
MESSAGE	6	7	8	10	12	13	16	20	22	25
MiniCAM Base	5	6	7	8	11	12	14	18	20	22
5th Scenario	5	6	6	8	10	11	14	17	19	21

Scenario	FUND									
IMAGE	-9	-5	-4	-1	2	3	6	10	14	24
MERGE Optimistic	-6	-4	-2	0	3	4	6	11	15	26
MESSAGE	-10	-6	-4	-1	1	2	5	9	12	21
MiniCAM Base	-7	-4	-2	0	3	4	6	11	14	25
5th Scenario	-11	-7	-5	-3	0	0	3	5	7	13

Table A5: Additional Summary Statistics of 2020 Global SC-CO₂ Estimates

Discount rate:	5.0%				3.0%				2.5%			
Statistic:	Mean	Variance	Skewness	Kurtosis	Mean	Variance	Skewness	Kurtosis	Mean	Variance	Skewness	Kurtosis
DICE	12	26	2	15	38	409	3	24	57	1097	3	30
PAGE	21	1481	5	32	68	13712	4	22	97	26878	4	23
FUND	3	41	5	179	19	1452	-42	8727	33	6154	-73	14931

Appendix B

The November 2013 revision of this TSD is based on two corrections to the runs based on the FUND model. First, the potential dry land loss in the algorithm that estimates regional coastal protections was misspecified in the model's computer code. This correction is covered in an erratum to Anthoff and Tol (2013a) published in the same journal (*Climatic Change*) in October 2013 (Anthoff and Tol (2013b)). Second, the equilibrium climate sensitivity distribution was inadvertently specified as a truncated Gamma distribution (the default in FUND) as opposed to the truncated Roe and Baker distribution as was intended. The truncated Gamma distribution used in the FUND runs had approximately the same mean and upper truncation point, but lower variance and faster decay of the upper tail, as compared to the intended specification based on the Roe and Baker distribution. The difference between the original estimates reported in the May 2013 version of this TSD and this revision are generally one dollar or less.

The July 2015 revision of this TSD is based on two corrections. First, the DICE model had been run up to 2300 rather than through 2300, as was intended, thereby leaving out the marginal damages in the last year of the time horizon. Second, due to an indexing error, the results from the PAGE model were in 2008 U.S. dollars rather than 2007 U.S. dollars, as was intended. In the current revision, all models have been run through 2300, and all estimates are in 2007 U.S. dollars. On average the revised SC-CO₂ estimates are one dollar less than the mean SC-CO₂ estimates reported in the November 2013 version of this TSD. The difference between the 95th percentile estimates with a 3% discount rate is slightly larger, as those estimates are heavily influenced by results from the PAGE model.

The July 2016 revision provides additional discussion of uncertainty in response to recommendations from the National Academy of Sciences, Engineering, and Medicine. It does not revisit the IWG's 2010 methodological decisions or update the schedule of SC-CO₂ estimates presented in the July 2015 revision. The IWG is currently seeking external expert advice from the National Academies on the technical merits and challenges of potential approaches to future updates of the SC-CO₂ estimates presented in this TSD. To date, the Academies' committee has issued an interim report that recommended against a near-term update to the SC-CO₂ estimates, but included recommendations for enhancing the presentation and discussion of uncertainty around the current estimates. This revision includes additional information that the IWG determined was appropriate to respond to these recommendations. Specifically, the executive summary presents more information about the range of quantified uncertainty in the SC-CO₂ estimates (including a graphical representation of symmetric high and low values from the frequency distribution of SC-CO₂ estimates conditional on each discount rate), and a new section has also been added that provides a unified discussion of the various sources of uncertainty and how they were handled in estimating the SC-CO₂. Efforts to make the sources of uncertainty clear have also been enhanced with the addition of a new appendix that describes in more detail the uncertain parameters in both the FUND and PAGE models (Appendix C). Furthermore, the full set of SC-CO₂ modeling results, which have previously been available upon request, are now provided on the OMB website for easy access. The Academies' final report (expected in early 2017) will provide longer term recommendations for a more comprehensive update. For more information on the status of the Academies' process, see: http://sites.nationalacademies.org/DBASSE/BECS/CurrentProjects/DBASSE_167526.

Appendix C

This appendix provides a general overview of the parameters that are treated probabilistically in each of the three integrated assessment models the IWG used to estimate the SC-CO₂. In the DICE model the only uncertain parameter considered was the equilibrium climate sensitivity as defined by the probability distribution harmonized across the three models. By default, all of the other parameters in the model are defined by point estimates and these definitions were maintained by the IWG. In the FUND and PAGE models many of the parameters, beyond the equilibrium climate sensitivity, are defined by probability distributions in the default versions of the models. The IWG maintained these default assumptions and allowed these parameters to vary in the Monte Carlo simulations conducted with the FUND and PAGE models.

Default Uncertainty Assumptions in FUND

In the version of the FUND model used by the IWG (version 3.8.1) over 90 of the over 150 parameters in the model are defined by probability distributions instead of point estimates, and for 30 of those parameters the values vary across the model's 16 regions. This includes parameters related to the physical and economic components of the model. The default assumptions in the model include parameters whose probability distributions are based on the normal, Gamma, and triangular distributions. In most cases the distributions are truncated from above or below. The choice of distributions and parameterizations are based on the model developers' assessment of the scientific and economic literature. Complete information on the exact probability distributions specified for each uncertain parameter is provided through the model's documentation, input data, and source code, available at: <http://www.fund-model.org/home>.

The physical components of the model map emissions to atmospheric concentrations, then map those concentrations to radiative forcing, which is then mapped to changes in global mean temperature. Changes in temperature are then used to estimate sea level rise. The parameters treated probabilistically in these relationships may be grouped into three main categories: atmospheric lifetimes, speed of temperature response, and sea level rise. First, atmospheric concentrations are determined by one box models, that capture a single representative sink, for each of the three non-CO₂ GHGs and a five box model for CO₂, that represents the multiple sinks in the carbon cycle that operate on different time frames. In each of these boxes, the lifetime of additions to the atmospheric concentration in the box are treated as uncertain. Second, parameters associated with speed at which the climate responds to changes in radiative forcing are treated as uncertain. In the FUND model radiative forcing, R_t , is mapped to changes in global mean temperature, T_t , through

$$T_t = T_{t-1} + \frac{1}{\theta_1 + \theta_2 ECS + \theta_3 ECS^2} \left(\frac{\psi ECS}{\ln(2)} R_t - T_{1-t} \right),$$

where the probability distribution for the equilibrium climate sensitivity, ECS , was harmonized across the models as discussed in the 2010 TSD. The parameters θ_i define the speed at which the temperature anomaly responds to changes in radiative forcing and are treated as uncertain in the model. Third, sea level rise is treated as a mean reverting function, where the mean is determined as proportional to the current global mean temperature anomaly. Both this proportionality parameter and the rate of mean reversion in this relationship are treated as uncertain in the model.

The economic components of the model map changes in the physical components to monetized damages. To place the uncertain parameters of the model associated with mapping physical endpoints to damages in context, it is useful to consider the general form of the damage functions in the model. Many of the damage functions in the model have forms that are roughly comparable to

$$D_{r,t} = \alpha_r Y_{r,t} \beta_{r,t} \left(\frac{y_{r,t}}{y_{r,b}} \right)^\gamma \left(\frac{N_{r,t}}{N_{r,b}} \right)^\phi T_t^\delta, \quad (1)$$

where α_r is the damage at a 1 °C global mean temperature increase as a fraction of regional GDP, $Y_{r,t}$. The model considers numerous changes that may reduce a region's benchmark vulnerability to climate change. For example, γ represents the elasticity of damages with respect to changes in the region's GDP per capita, $y_{r,t}$, relative to a benchmark value, $y_{r,b}$; ϕ represents the elasticity of damages with respect to changes in the region's population, $N_{r,t}$, relative to a benchmark value, $N_{r,b}$; and the projection $\beta_{r,t}$ provides for an exogenous reduction in vulnerability (e.g., forecast energy efficiency improvements that affect space cooling costs). Once the benchmark damages have been scaled due to changes in vulnerability they are adjusted based on a non-linear scaling of the level of climate change forecast, using a power function with the exponent, δ .

Some damage categories have damage function specifications that differ from the example in (1). For example, agriculture and forestry damages take atmospheric concentrations of CO₂ and the rate of climate change into account in different forms, though the method by which they calculate the monetized impact in these cases is similar with respect to accounting for GDP growth and changes in vulnerability. In other cases the process by which damages are estimated is more complex. For example, in estimating damages from sea level rise the model considers explicit regional decision makers that choose levels of coastal protection in a given year based on a benefit-cost test. In estimating the damages from changes in cardiovascular mortality risk the model considers forecast changes in the proportion of the population over the age of 65 and deemed most vulnerable by the model developers. Other damage categories may also have functional forms that differ slightly from (1), but in general this form provides a useful framework for discussing the parameters for which the model developers have defined probability distributions as opposed to point estimates.

In many damage categories (e.g., sea level rise, water resources, biodiversity loss, agriculture and forestry, and space conditioning) the benchmark damages, α_r , are treated as uncertain parameters in the model and in most case they are assumed to vary by region. The elasticity of damages with respect to changes in regional GDP per capita, γ , and the elasticity with respect to changes in regional population, ϕ , are also treated as uncertain parameters in most damage functions in the model, though they are not assumed to vary across regions. In most cases the exponent, δ , on the power function that scales damages based on the forecast level of climate change are also treated as uncertain parameters, though they are not assumed to vary across regions in most cases.

Figure C1 presents results of an analysis from the developers of the FUND model that examines the uncertain parameters that have the greatest influence on estimates of the SC-CO₂ based on the default version of the model. While some of the modeling inputs are different for the SC-CO₂ estimates calculated by the IWG these parameters are likely to remain highly influential in the FUND modeling results.




Figure C1: Influence of Key Uncertain Parameters in Default FUND Model (Anthoff and Tol 2013a)¹⁹

Default Uncertainty Assumptions in PAGE

In the version of the PAGE model used by the IWG (version PAGE09) there are over 40 parameters defined by probability distributions instead of point estimates.²⁰ The parameters can broadly be classified as related to climate science, damages, discontinuities, and adaptive and preventive costs. In the default version of the model, all of the parameters are modeled as triangular distributions except for the one variable related to the probability of a discontinuity occurring, which is represented by a uniform distribution. More detail on the model equations can be found in Hope (2006, 2011a) and the default minimum, mode, and maximum values for the parameters are provided in Appendix 2 of Hope (2011a). The calibration of these distributions is based on the developer's assessment of the IPCC's Fourth Assessment report and scientific articles referenced in Hope (2011a, 2011b, 2011c). The IWG added an uncertain parameter to the default model, specifically the equilibrium climate sensitivity parameter, which was harmonized across the models as discussed in the 2010 TSD.

In the climate component of the PAGE model, atmospheric CO₂ concentration is assumed to follow an initial rapid decay followed by an exponential decline to an equilibrium level. The parameters treated probabilistically in this decay are the proportion of the anthropogenic CO₂ emissions that enter the atmosphere, the half-life of the CO₂'s atmospheric residence, and the fraction of cumulative emissions that ultimately remains in the atmosphere. A carbon cycle feedback is included to represent the impact of increasing temperatures on the role of the terrestrial biosphere and oceans in the carbon cycle. This feedback is modeled with probabilistic parameters representing the percentage increase in the CO₂ concentration anomaly and with an uncertain upper bound on this percentage.

The negative radiative forcing effect from sulfates is modeled with probabilistic parameters for the direct linear effect due to backscattering and the indirect logarithmic effect assumed for cloud interactions. The radiative forcing from CO₂, all other greenhouse gases, and sulfates are combined in a one box model to estimate the global mean temperature. Uncertainty in the global mean temperature response to change in radiative forcing is based on the uncertain equilibrium climate sensitivity parameter and uncertainty in the half-life of the global response to an increase in radiative forcing, which defines the inertia of the climate system in the model. Temperature anomalies in the model vary geographically, with larger increases over land and the poles. Probabilistic parameters are used for the ratios of the temperature anomaly over land relative to the ocean and the ratio of the temperature anomaly over the poles relative to the equator. The PAGE model also includes an explicit sea level component, modelled as a lagged function of the global mean temperature anomaly. The elements of this component that are treated

¹⁹ Based on a coefficients of standardized regression of parameter draws on the SC-CO₂ using FUND 3.8.1 under Ramsey discounting with a pure rate of time preference of one percent and rate of relative risk aversion of 1.5. The 90 percent confidence intervals around the regression coefficients are presented as error bars.

²⁰ This appendix focuses on the parameters in the PAGE model related to estimating the climate impacts and principle calculation of the monetized damages. There are over 60 additional parameters in the model related to abatement and adaptation, which may be highly relevant for purposes other than estimating the SC-CO₂, but are not discussed here.

probabilistically include: sea level rise from preindustrial levels to levels in the year 2000, the asymptotic sea level rise expected with no temperature change, the predicted sea level rise experience with a temperature change, and the half-life of the sea level rise.

In the economic impacts module, damages are estimated for four categories: sea level rise, economic damages, non-economic damages, and damages from a discontinuity. Each damage category is calculated as a loss proportional to GDP. The model first calculates damages for a “focus region” (set to the European Union) assuming the region’s base year GDP per capita. Damages for other regions are assumed to be proportional to the focus region’s damage, represented by a regional weighting factor.

Economic damages, non-economic damages, and damages from sea level rise are modeled as polynomial functions of the temperature or sea level impact, which are defined as the regional temperature or sea level rise above a regional tolerable level. These functions are calibrated to damages at some reference level (e.g., damages at 3°C or damages for a ½ meter sea level rise). The specification allows for the possibility of “initial benefits” from small increases in regional temperature. The variables represented by a probability distributions in this specification are: the regional weighting factors; the initial benefits; the calibration point; the damages at the calibration point; and the exponent on the damage functions.

The damages from a discontinuity are treated differently from other damages in PAGE because the event either occurs or it does not in a given model simulation. In the PAGE model, the probability of a discontinuity is treated as a discrete event, where if it occurs, additional damages would be borne and therefore added to the other estimates of climate damages. Uncertain parameters related to this discontinuity include the threshold global mean temperature beyond which a discontinuity becomes possible and the increase in the probability of a discontinuity as the temperature anomaly continues to increase beyond this threshold. If the global mean temperature has exceeded the threshold for any time period in a model run, then the probability of a discontinuity occurring is assigned, otherwise the probability is set to zero. For each time period a uniform random variable is drawn and compared to this probability to determine if a discontinuity event has occurred in that simulation. The additional loss if a discontinuity does occur in a simulation is represented by an uncertain parameter and is multiplied by the uncertain regional weighting factor to obtain the regional effects.

Damages for each category in each region are adjusted to account for the region’s forecast GDP in a given model year to reflect differences in vulnerability based on the relative level of economic development. Specifically, the damage estimates are multiplied by a factor equal to the ratio of a region’s actual GDP per capita to the base year GDP per capita, where the ratio exponentiated with a value less than or equal to zero. The exponents vary across damage categories and in each case are treated as uncertain parameters.

Finally, in each region damages for each category are calculated sequentially (sea level rise, economic, non-economic, and discontinuity, in that order) and are assessed to ensure that they do not create total damages that exceed 100 percent of GDP for that region. Damages transition from a polynomial function to a logistic path once they exceed a certain proportion of remaining GDP, and the proportion where this transition begins is treated as uncertain. An additional parameter labeled the “statistical value of

civilization,” also treated as uncertain, caps total damages (including abatement and adaptation costs described below) at some maximum level.

Figure C2 presents results of an analysis from the developers of the PAGE model that examines the uncertain parameters that have the greatest influence on estimates of the SC-CO₂ based on the default version of the model. Although some of the modeling inputs are different for the SC-CO₂ estimates calculated by the IWG, these parameters are likely to remain highly influential in the PAGE modeling results.




Figure C2: Influence of Key Uncertain Parameters in Default PAGE Model (Hope 2013)²¹

²¹ Based on a standardized regression of the parameters. The values give the predicted increase in the SC-CO₂ in 2010 based on a one standard deviation increase in the coefficient, using the default parameters for PAGE09 under Ramsey discounting with an uncertain pure rate of time preference and rate of relative risk aversion.